

Perioperative Management of Patients With Atrial Fibrillation Receiving a Direct Oral Anticoagulant

James D. Douketis, MD; Alex C. Spyropoulos, MD; Joanne Duncan, BSc; Marc Carrier, MD, MSc; Gregoire Le Gal, MD; Alfonso J. Tafur, MD; Thomas Vanassche, MD; Peter Verhamme, MD; Sudeep Shivakumar, MD; Peter L. Gross, MD, MSc; Agnes Y. Y. Lee, MD, MSc; Erik Yeo, MD; Susan Solymoss, MD; Jeannine Kassis, MD; Geneviève Le Templier, MD; Stephen Kowalski, MD; Mark Blostein, MD; Vinay Shah, MD; Elizabeth MacKay, MD; Cynthia Wu, MD; Nathan P. Clark, PharmD; Shannon M. Bates, MDCM, MSc; Frederick A. Spencer, MD; Eleni Arnaoutoglou, MD, PhD; Michiel Coppens, MD, PhD; Donald M. Arnold, MD, MSc; Joseph A. Caprini, MD; Na Li, PhD; Karen A. Moffat, MLT; Summer Syed, MD, MSc; Sam Schulman, MD, PhD

[+ Supplemental content](#)

IMPORTANCE Patients with atrial fibrillation (AF) who use a direct oral anticoagulant (DOAC) and request elective surgery or procedure present a common clinical situation yet perioperative management is uncertain.

OBJECTIVE To investigate the safety of a standardized perioperative DOAC management strategy.

DESIGN, SETTING, AND PARTICIPANTS The Perioperative Anticoagulation Use for Surgery Evaluation (PAUSE) cohort study conducted at 23 clinical centers in Canada, the United States, and Europe enrolled and screened patients from August 1, 2014, through July 31, 2018. Participants (n = 3007) had AF; were 18 years of age or older; were long-term users of apixaban, dabigatran etexilate, or rivaroxaban; were scheduled for an elective surgery or procedure; and could adhere to the DOAC therapy interruption protocol.

INTERVENTIONS A simple standardized perioperative DOAC therapy interruption and resumption strategy based on DOAC pharmacokinetic properties, procedure-associated bleeding risk, and creatinine clearance levels. The DOAC regimens were omitted for 1 day before a low-bleeding-risk procedure and 2 days before a high-bleeding-risk procedure. The DOAC regimens were resumed 1 day after a low-bleeding-risk procedure and 2 to 3 days after a high-bleeding-risk procedure. Follow-up of patients occurred for 30 days after the operation.

MAIN OUTCOMES AND MEASURES Major bleeding and arterial thromboembolism (ischemic stroke, systemic embolism, and transient ischemic attack) and the proportion of patients with an undetectable or minimal residual anticoagulant level (<50 ng/mL) at the time of the procedure.

RESULTS The 3007 patients with AF (mean [SD] age of 72.5 [9.39] years; 1988 men [66.1%]) comprised 1257 (41.8%) in the apixaban cohort, 668 (22.2%) in the dabigatran cohort, and 1082 (36.0%) in the rivaroxaban cohort; 1007 patients (33.5%) had a high-bleeding-risk procedure. The 30-day postoperative rate of major bleeding was 1.35% (95% CI, 0%-2.00%) in the apixaban cohort, 0.90% (95% CI, 0%-1.73%) in the dabigatran cohort, and 1.85% (95% CI, 0%-2.65%) in the rivaroxaban cohort. The rate of arterial thromboembolism was 0.16% (95% CI, 0%-0.48%) in the apixaban cohort, 0.60% (95% CI, 0%-1.33%) in the dabigatran cohort, and 0.37% (95% CI, 0%-0.82%) in the rivaroxaban cohort. In patients with a high-bleeding-risk procedure, the rates of major bleeding were 2.96% (95% CI, 0%-4.68%) in the apixaban cohort and 2.95% (95% CI, 0%-4.76%) in the rivaroxaban cohort.

CONCLUSIONS AND RELEVANCE In this study, patients with AF who had DOAC therapy interruption for elective surgery or procedure, a perioperative management strategy without heparin bridging or coagulation function testing was associated with low rates of major bleeding and arterial thromboembolism.

JAMA Intern Med. doi:10.1001/jamainternmed.2019.2431
Published online August 5, 2019.

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: James D. Douketis, MD, Department of Medicine, McMaster University and St Joseph's Healthcare Hamilton, 50 Charlton Ave E, Room F-544, Hamilton, Ontario L8N 4A6, Canada (jdouket@mcmaster.ca).

The perioperative management of patients who receive a direct oral anticoagulant (DOAC) for atrial fibrillation (AF) and require elective surgery or procedure is a common clinical scenario for which best practices are uncertain.¹ Each year, 1 in 6 patients with AF, or an estimated 6 million patients worldwide, will require perioperative anticoagulant management.^{2,3} When DOAC regimens became available for clinical use in AF, starting in 2010, no studies had been conducted to inform the timing of perioperative DOAC therapy interruption and resumption, whether heparin bridging should be given, and whether preoperative coagulation function testing was needed.⁴ Uncertainty about the perioperative management of DOACs may be associated with unsubstantiated practices and increased harm to patients. Thus, a DOAC therapy interruption interval that is too long may increase the risk for thromboembolism, whereas an interruption interval that is too short may increase the risk for bleeding which, in turn, delays anticoagulant resumption.⁵ Perioperative heparin bridging has been used in DOAC-treated patients,⁶ but this practice does not make pharmacologic sense given the short, 8- to 14-hour DOAC elimination half-lives,⁴ its association with increased bleeding, and its questionable efficacy.^{6,7} Preoperative coagulation testing has been suggested to identify patients with an excessive residual anticoagulant level in whom a procedure can be delayed or the DOAC reversed,⁸ but this suggestion is problematic because DOAC-specific coagulation tests are not widely available, reference ranges are lacking, and such testing may not be advantageous for patients.⁹

Most clinical studies investigating perioperative DOAC regimen management are retrospective subanalyses of randomized clinical trials that assessed DOAC regimens for stroke prevention in AF or are patient registries.^{3,10-13} One study that assessed standardized perioperative management included only patients who were taking dabigatran etexilate.¹⁴ The perioperative management of DOAC regimens varies widely in clinical practice,¹⁵ and practice guidelines provide weak and inconsistent management recommendations.¹⁶⁻²⁰

We designed the Perioperative Anticoagulation Use for Surgery Evaluation (PAUSE) study to assess the safety of a standardized perioperative management strategy for a DOAC regimen. We hypothesized that a simple management approach, which is based on DOAC-specific interruption and resumption intervals, forgoes perioperative heparin bridging, and does not require preoperative coagulation function testing, is safe to use for patient care. For each DOAC cohort that received DOAC-specific perioperative management, we defined safety as excluding 30-day perioperative rates of major bleeding of 2% and arterial thromboembolism of 1.5%, according to expected outcome rates (1% for major bleeding and 0.5% for arterial thromboembolism) observed with optimal perioperative management of warfarin sodium^{3,21} and with a proof-of-concept prospective study of standardized perioperative dabigatran management.¹⁴ We also postulated that this management would yield a high proportion of patients (>90%) with an undetectable or minimal residual anticoagulant level at the time of the procedure.

Key Points

Question Is a standardized perioperative management approach safe for patients with atrial fibrillation who use a direct oral anticoagulant and require elective surgery or procedure?

Findings In this cohort study of 3007 patients with atrial fibrillation using apixaban, dabigatran, or rivaroxaban, the direct oral anticoagulant treatment was stopped and resumed before and/or after elective surgery or procedure using standardized protocols without heparin bridging. The 30-day postoperative rates of major bleeding were less than 2%, and the rates of stroke were less than 1%.

Meaning In this study, in patients treated with a direct oral anticoagulant, a simple standardized perioperative management approach was associated with low rates of bleeding and stroke.

Methods

Study Design and Oversight

The PAUSE study design and data analysis plan were developed by the steering committee and are described elsewhere.²² The study was managed by the McMaster Centre for Transfusion Research, which was responsible for the study organization as well as data collection, validation, maintenance, and analysis. Study data were collected and managed using REDCap electronic data capture tools.²³ The institutional review board of each of the 23 participating clinical center in Canada, the United States, and Europe approved PAUSE, and all study participants provided written informed consent.

PAUSE is a prospective management study involving DOAC-treated patients with AF who required anticoagulant therapy interruption for elective surgery or procedure. Patients were separated into 3 cohorts on the basis of DOAC used (apixaban, dabigatran, or rivaroxaban) and received standardized perioperative management according to the DOAC. A randomized clinical trial design was considered to assess the proposed (experimental) management but was not adopted because no alternative strategy existed that would be suitable as a comparator (control) management. For example, a management approach that omits DOAC regimens for a longer (4- to 6-day) preoperative period, as suggested in other studies,¹⁶ would not make clinical sense as a comparator given the short DOAC elimination half-lives; moreover, the longer period without anticoagulation might expose patients to an increased thromboembolic risk. Similarly, adopting unspecified usual care as a comparator would also be unsuitable, as the usual care may be too heterogeneous to allow a meaningful comparison to the standardized, more uniform management used in this study.¹⁵ A cohort design is appropriate for assessing a management strategy when expected rates of clinical outcomes are low (0.5%-1% in PAUSE) and when there is sufficient statistical power to exclude clinically important higher outcome rates (1.5%-2% in PAUSE).^{23,24}

Patients

Consecutive patients with the following characteristics were assessed for study eligibility: adults (aged ≥18 years) with AF

Figure. Perioperative Direct Oral Anticoagulant (DOAC) Management Protocol

DOAC	Surgical Procedure-Associated Bleeding Risk	Preoperative DOAC Interruption Schedule					Day of Surgical Procedure (No DOAC)	Postoperative DOAC Resumption Schedule			
		Day -5	Day -4	Day -3	Day -2	Day -1		Day +1	Day +2	Day +3	Day +4
Apixaban	High	[Orange arrow from Day -5 to Day -2, thickened from Day -4 to Day -2]					[Vertical bar]	[Orange arrow from Day +2 to Day +4, thickened from Day +2 to Day +3]			
	Low	[Dark blue arrow from Day -5 to Day -1]						[Dark blue arrow from Day +1 to Day +4]			
Dabigatran etexilate (CrCl ≥50 mL/min)	High	[Orange arrow from Day -5 to Day -2, thickened from Day -4 to Day -2]						[Orange arrow from Day +2 to Day +4, thickened from Day +2 to Day +3]			
	Low	[Dark blue arrow from Day -5 to Day -1]						[Dark blue arrow from Day +1 to Day +4]			
Dabigatran etexilate (CrCl <50 mL/min) ^a	High	[Light blue arrow from Day -5 to Day -4]	[Shaded]					[Orange arrow from Day +2 to Day +4, thickened from Day +2 to Day +3]			
	Low	[Dark blue arrow from Day -5 to Day -1]						[Dark blue arrow from Day +1 to Day +4]			
Rivaroxaban	High	[Orange arrow from Day -5 to Day -2, thickened from Day -4 to Day -2]						[Orange arrow from Day +2 to Day +4, thickened from Day +2 to Day +3]			
	Low	[Dark blue arrow from Day -5 to Day -1]						[Dark blue arrow from Day +1 to Day +4]			

No DOAC was taken on certain days (shaded) and on the day of the elective surgery or procedure. The light blue arrows refer to an exception to the basic management, a subgroup of patients taking dabigatran with a creatinine clearance (CrCl) less than 50 mL/min. The orange arrows refer to patients having a high-bleed-risk surgical procedure. Dark blue arrows refer to patients having a

low-bleed-risk surgical procedure. The thickened orange part of arrows refer to flexibility in the timing of DOAC resumption after a procedure.

^a Cancer diagnosed within 3 months or has been treated within 6 months or metastatic.

who were long-term users of apixaban (5 mg or 2.5 mg twice daily), dabigatran etexilate (150 mg or 110 mg twice daily), or rivaroxaban (20 mg or 15 mg daily); were scheduled to have an elective surgery or procedure that required interruption of the anticoagulant regimen; and were able to adhere to the DOAC therapy interruption protocol at the time of enrollment. Patients were excluded if they fit 1 or more of the following criteria: creatinine clearance (CrCl) level less than 25 mL/min for apixaban users or CrCl level less than 30 mL/min for dabigatran or rivaroxaban users (to convert CrCl level to milliliters per second per meter squared, multiply by 0.0167),²⁵ cognitive impairment or psychiatric illness, did not consent to participate, previous study participation, or more than 1 procedure planned within 30 days. Before the procedure, patients were categorized as having a high- or low-bleeding-risk procedure according to a prespecified classification (eAppendix 1 in the Supplement)⁷; this classification informed the timing of DOAC therapy interruption and resumption.²² Our aim was that at least one-third of patients enrolled into each DOAC cohort would be classified as high bleeding risk.

Procedures

The perioperative management strategy for a DOAC regimen was designed with 2 broad aims: (1) to have the shortest duration of DOAC therapy interruption before and after the procedure so as to minimize the risks for bleeding and thromboembolism, and (2) to have a simple interruption and resumption protocol for each DOAC that would be easy to use by clinicians and easily understood by patients.

Patients were enrolled and managed using a standardized perioperative DOAC strategy based on DOAC pharmacokinetic properties (10- to 14-hour half-lives, and 1- to 3-hour peak action), the procedure-associated bleeding risk, and patient CrCl level (Figure).²² Before the procedure, DOAC regimens were omitted for 1 day before a low-bleeding-risk pro-

cedure (36- to 42-hour interval corresponding to approximately 3 DOAC half-lives) and were omitted 2 days before a high-bleeding-risk procedure (60- to 68-hour interval corresponding to approximately 5 DOAC half-lives). Patients using dabigatran with a CrCl level less than 50 mL/min had longer interruption intervals to account for renal dependence of dabigatran clearance.¹ Blood samples were taken from patients just before the procedure to measure their residual anticoagulant level, but these results were not available for clinical use. Plasma samples were frozen and stored at each clinical site and later analyzed in a centralized laboratory using standardized blood processing and assay methods (eAppendix 2 in the Supplement). After the operation, DOAC regimens were resumed 1 day (approximately 24 hours) after a low-bleeding-risk procedure and 2 to 3 days (48-72 hours) after a high-bleeding-risk procedure, provided that hemostasis was achieved. Patient thromboembolic risk, based on the CHADS₂ (congestive heart failure, hypertension, aged 75 years or older, diabetes, and previous stroke or transient ischemic attack) risk score, did not affect perioperative DOAC regimen management because this risk score is used in a perioperative setting to assess the need for heparin bridging, which was not performed in the present study.^{26,27} Patients at high risk for venous thromboembolism could receive a prophylactic dose of heparin after the operation until DOAC therapy resumption.

Clinical Outcomes and Residual Anticoagulant Level

Study clinical outcomes were assessed from the time the first DOAC dose was interrupted until 30 days after the operation. Patients had scheduled weekly telephone follow-up and additional clinic visits as needed to document clinical outcomes. The primary clinical outcomes were major bleeding and arterial thromboembolism (ischemic stroke, transient ischemic attack, and systemic embolism). The secondary clinical outcomes were clinically relevant nonmajor bleeding, minor

bleeding, death, myocardial infarction, deep vein thrombosis, pulmonary embolism, and catheter-associated venous or arterial thrombosis. Study outcomes were defined according to standardized criteria (eAppendix 3 in the [Supplement](#))^{28,29} and were independently adjudicated by a committee that was blinded to the DOAC cohort, procedure bleeding risk, and preoperative DOAC treatment levels.

The residual anticoagulant level just before the procedure was measured by DOAC-specific anti-factor Xa assays for apixaban and rivaroxaban as well as by the dilute thrombin time for dabigatran.³⁰ The residual anticoagulant level was also measured with nonspecific coagulation tests: prothrombin time, international normalized ratio, activated partial thromboplastin time, and thrombin time.³¹

Study Hypothesis and Sample Size Determination

We hypothesized that, for each DOAC cohort, the PAUSE management would be associated with a 1% rate of major bleeding (with the upper limit of the 1-sided 95% CI to exclude a 2% rate) and a 0.5% rate of arterial thromboembolism (with the upper limit of the 1-sided 95% CI to exclude a 1.5% rate). Thus, the null hypothesis was that the proposed protocol was unsafe; that is, the proportion of the major bleeding (or arterial thromboembolism) was 2% or higher (or $\geq 1.5\%$); an alternative hypothesis was that the protocol was safe; that is, the proportion was lower than 2% (or $< 1.5\%$). A 1-sided $P < .05$ was considered statistically significant, and a statistically significant result would mean that, with the 1-sided 95% CI, the true incidence of major bleeding was lower than 2% and arterial thromboembolism was lower than 1.5% for each DOAC cohort, rejecting the null hypothesis.

When PAUSE was designed in 2013, we were more confident about estimates, based on findings from available studies,^{3,21,32} of perioperative rates of major bleeding than arterial thromboembolism. Therefore, major bleeding was the primary determinant of sample size, and the sample size calculation was based on an expected rate of 1%. The required sample size was 987 patients per DOAC cohort, which provided 80% power at the .05 significance level (1-sided) to detect a proportion that was lower than 2% for major bleeding. With this sample size, there was also 80% power at the 5% significance level (1-sided) to detect a proportion that was lower than 1.5% for arterial thromboembolism, based on an expected rate of 0.5%.

The number of patients per DOAC cohort was increased by 10% (to 1097) to anticipate cancelled operations and patients lost to follow-up. We also postulated that the DOAC therapy interruption protocol would yield more than 90% of patients with a preoperative residual anticoagulant level less than 50 ng/mL, which was considered empirically as a level that would allow a procedure to proceed safely.

Statistical Analysis

For the primary clinical outcomes, a 1-sided test for 1 proportion with continuity correction was used to determine within each DOAC cohort at the patient level if the proportion of major bleeding was lower than 2% and if the proportion of arterial thromboembolism was lower than 1.5%.³³ The primary

analysis was conducted in the main study population of patients who had at least 1 DOAC dose interrupted. For each primary outcome within each DOAC cohort, we reported the proportion and associated 1-sided 95% CI as well as the P value from the 1-sided test for 1 proportion to check that the outcome rate was lower than the expected rate of 2% for major bleeding and 1.5% for arterial thromboembolism. For the secondary clinical outcomes, we assessed rates of mortality and other adverse events for patients within each DOAC cohort. We reported the proportions with 2-sided 95% CIs of the secondary outcomes for each cohort.

For the preoperative residual anticoagulant level outcome, we identified the proportion of patients with an anti-factor Xa level (for apixaban or rivaroxaban) or dilute thrombin time (for dabigatran) of less than 50 ng/mL (30-49.9 ng/mL and < 30 ng/mL) or 50 ng/mL or greater; this calculation was done separately for patients with a low-bleeding risk and patients with a high-bleeding-risk procedure because the bleeding risk determined the DOAC therapy interruption interval, which would affect the residual anticoagulant level. We also identified the median (interquartile range [IQR]) prothrombin time, international normalized ratio, activated partial thromboplastin time, and thrombin time as well as the proportion of patients with an elevated prothrombin time, international normalized ratio, activated partial thromboplastin time, and thrombin time.

Because the analyses of secondary clinical outcomes and coagulation test outcomes were descriptive, no statistical hypothesis testing was considered. In this analysis, we assessed rates of major bleeding in patients according to procedure-associated bleeding risk as perioperative management differed between patients considered at high and low risk for bleeding. We also assessed rates of primary outcomes in a population of patients within each cohort who adhered to the DOAC therapy interruption and resumption protocols.

Results

Patients

We screened 3640 patients from August 1, 2014, through July 31, 2018, from 23 clinical sites in Canada, the United States, and Europe (eAppendix 4 in the [Supplement](#)). Of these patients, 3007 (82.6%) were enrolled and were included in the primary analysis: 1257 (41.8%) in the apixaban cohort, 668 (22.2%) in the dabigatran cohort, and 1082 (36.0%) in the rivaroxaban cohort (eFigure in the [Supplement](#)). The baseline characteristics of the patients in each DOAC cohort are shown in [Table 1](#). Overall, patients had a mean (SD) age of 72.5 (9.39) years and were predominantly male (1988 [66.1%]). The types of procedures that patients underwent in each DOAC cohort are shown in eAppendix 5 in the [Supplement](#).

Perioperative Anticoagulant Management

[Table 2](#) shows the DOAC therapy interruption intervals for the apixaban, dabigatran (≥ 50 mL/min and < 50 mL/min subgroups), and rivaroxaban cohorts as well as the DOAC therapy resumption intervals for the apixaban, dabigatran, and riva-

Table 1. Baseline Patients Characteristics

Variable	No. (%)		
	Apixaban Cohort (n = 1257)	Dabigatran Cohort (n = 668)	Rivaroxaban Cohort (n = 1082)
Age, mean (SD), y	73.1 (9.15)	72.4 (9.9)	72.0 (9.3)
Male	805 (64.0)	458 (68.6)	725 (67.0)
BMI, mean (SD)	29.49 (6.2)	30.24 (6.8)	29.8 (6.5)
Race/ethnicity			
White	1204 (95.8)	654 (97.9)	1045 (96.6)
Non-white	43 (3.4)	12 (1.8)	25 (2.3)
Unknown	10 (0.8)	2 (0.3)	12 (1.1)
Risk stratification scores, mean (SD)			
CHADS ₂ ^a	2.1 (1.3)	2.2 (1.3)	2.0 (1.3)
CHADS ₂ -VA ₂ Sc ^b	3.5 (1.7)	3.5 (1.6)	3.3 (1.6)
Modified HAS-BLED ^c	2.0 (0.9)	1.9 (0.9)	1.8 (0.9)
Medical condition			
Congestive heart failure	243 (19.3)	111 (16.6)	140 (12.9)
Hypertension	933 (74.2)	504 (75.4)	784 (72.5)
Diabetes	337 (26.8)	185 (27.7)	273 (25.2)
Stroke	98 (7.8)	64 (9.6)	77 (7.1)
Transient ischemic attack	117 (9.3)	93 (13.9)	99 (9.1)
Coronary artery disease	232 (18.5)	113 (16.9)	177 (16.4)
Peripheral arterial disease	8 (0.6)	6 (0.9)	13 (1.2)
Bioprosthetic heart valve	35 (2.8)	10 (1.5)	20 (1.8)
Mitral valve disease	125 (9.9)	51 (7.6)	86 (7.9)
Venous thromboembolism	77 (6.1)	40 (6.0)	85 (7.9)
Active cancer ^d	105 (8.3)	57 (8.5)	107 (9.9)
Laboratory values, mean (SD)			
Hemoglobin, g/L	134.4 (17.8)	140.1 (50.0)	136.8 (31.6)
Platelets <100 × 10 ⁶ /L	8 (0.6)	2 (0.3)	3 (0.3)
Serum creatinine, μmol/L	94.1 (28.8)	87.7 (21.6)	90.3 (22.5)
Creatinine clearance, ml/min ^e	77.9 (32.0)	85.9 (35.7)	82.2 (32.8)
Medication use			
Lower-dose DOAC regimen ^f	252 (20.0)	248 (37.1)	181 (16.7)
Aspirin	156 (12.4)	98 (14.7)	99 (9.1)
P2Y ₁₂ inhibitor ^g	12 (0.9)	7 (1.0)	11 (1.0)
P-glycoprotein or cytochrome P450 3A4 inhibitor or inducer ^h	76 (6.0)	53 (7.9)	55 (5.1)
Elective surgery or procedure type			
High bleeding risk	406 (32.3)	228 (34.1)	373 (34.5)
Low bleeding risk	851 (67.7)	440 (65.9)	709 (65.5)
Anesthesia type			
General	410 (32.6)	193 (28.9)	384 (35.5)
Neuraxial	103 (8.2)	57 (8.5)	70 (6.5)
Other	689 (54.8)	369 (55.2)	584 (54.0)

Abbreviations: BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); DOAC, direct oral anticoagulant.

SI conversion factor: To convert creatinine clearance to milliliters per second per square meter, multiply by 0.0167.

^a CHADS₂ risk score range: 1-6; risks include congestive heart failure, hypertension, age 75 years or older, diabetes, and previous stroke or transient ischemic attack.

^b CHADS₂-VA₂Sc risk score range: 1-9; risks include congestive heart failure, hypertension, age 75 years or older or 65 years or older, diabetes, previous stroke or transient ischemic attack, female sex, and vascular disease.

^c HAS-BLED bleeding risk score range: 1-7; risks include hypertension, abnormal renal or liver function, previous stroke, previous bleed or bleed predisposition, labile international normalized ratio (omitted), age 65 years or older, and drug use that affects hemostasis or alcohol use (omitted).

^d Cancer diagnosed within 3 months or treated within 6 months or metastatic.

^e Based on Cockcroft-Gault formula.

^f Apixaban 2.5 mg twice daily, or dabigatran etexilate 110 mg twice daily, or rivaroxaban 15 mg daily.

^g Clopidogrel bisulfate, ticagrelor, prasugrel hydrochloride, or ticlopidine hydrochloride.

^h Drugs that can inhibit or induce DOAC activity (eAppendix 9 in the Supplement).

roxaban cohorts. Of the 3007 patients in the primary analysis cohort (≥1 dose interrupted), 159 (5.3%) deviated from the DOAC therapy interruption protocol, 202 (6.7%) deviated from the DOAC therapy resumption protocol, and 22 (0.7%) were lost to follow-up, leaving 2624 patients (87.3%) to be included in the per protocol analysis.

Study Outcomes

In the primary analysis cohort (Table 3), the 30-day postoperative rate of major bleeding was 1.35% (95% CI, 0%-2.00%)

in the apixaban cohort, 0.90% (95% CI, 0%-1.73%) in the dabigatran cohort, and 1.85% (95% CI, 0%-2.65%) in the rivaroxaban cohort. The rate of arterial thromboembolism was 0.16% (95% CI, 0%-0.48%) in the apixaban cohort, 0.60% (95% CI, 0%-1.33%) in the dabigatran cohort, and 0.37% (95% CI, 0%-0.82%) in the rivaroxaban cohort. All 43 major bleeding events occurred postoperatively at a median (IQR) of 2 (0-6) days; 9 of 10 arterial thromboembolic events occurred postoperatively at a median (IQR) of 2 (0-6) days. Rates of major bleeding according to procedure-associated bleeding risk are

Table 2. Preoperative and Postoperative Direct Oral Anticoagulant Interruption and Resumption

Cohort	Preoperative Management			Postoperative Management			Patient Adherence to Resumption Protocol, No. (%)	Patient Receipt of Prophylactic-Dose LMWH, No. (%)
	DOAC Preoperative Omission, No. (IQR), d	Interruption Interval (IQR), h	Interruption Protocol, No. (%)	DOAC Postoperative Resumption, No. (IQR), d	Resumption Interval (IQR), h	Resumption Protocol, No. (%)		
Apixaban								
Low bleeding risk (n = 851)	1 (1-1)	39.3 (37.4-41.5)	819 (96.24)	1 (1-1)	22.2 (19.3-31.9)	745 (87.5)	16 (1.9)	
High bleeding risk (n = 406)	2 (2-2)	63.8 (61-67)	378 (93.1)	3 (2-4)	67.8 (45.1-91.4)	399 (98.3)	133 (32.8)	
Dabigatran etexilate, CrCl ≥50 mL/min								
Low bleeding risk (n = 386)	1 (1-1)	39.7 (38-41.9)	368 (95.34)	NA	NA	NA	NA	
High bleeding risk (n = 202)	2 (2-2)	63.2 (61.5-67.2)	187 (92.57)	NA	NA	NA	NA	
Dabigatran, CrCl <50 mL/min								
Low bleeding risk (n = 54)	2 (2-2)	64.4 (62-66)	50 (92.59)	NA	NA	NA	NA	
High bleeding risk (n = 26)	4 (4-4)	110.2 (108.3-112.7)	22 (84.62)	NA	NA	NA	NA	
Dabigatran (all patients) ^a								
Low bleeding risk (n = 440)	NA	NA	NA	1 (1-1)	23 (20.5-33.6)	425 (96.6)	7 (1.6)	
High bleeding risk (n = 228)	NA	NA	NA	3 (2-3)	66.4 (45.1-81.4)	227 (99.6)	85 (37.3)	
Rivaroxaban								
Low bleeding risk (n = 709)	1 (1-1)	48 (40.7-51)	674 (95.06)	1 (1-1)	25 (20.8-33.5)	641 (90.41)	8 (1.13)	
High bleeding risk (n = 373)	2 (2-2)	72 (65.6-75)	350 (93.83)	3 (2-4)	69.4 (46.4-94)	370 (99.2)	131 (35.1)	

Abbreviations: CrCl, creatinine clearance; DOAC, direct oral anticoagulant; IQR, interquartile range; LMWH, low-molecular-weight heparin; NA, not applicable.

^a Postoperative resumption of dabigatran was the same for patients with a CrCl 50 milliliters per minute or more and less than 50 milliliters per minute.

SI conversion factor: To convert creatinine clearance to milliliters per second per square meter, multiply by 0.0167.

shown in Table 4; in the high-bleed-risk subgroups, the rate of major bleeding was 2.96% (95% CI, 0%-4.68%) in the apixaban cohort, 0.88 (95% CI, 0%-2.62%) in the dabigatran cohort, and 2.95% (95% CI, 0%-4.76%) in the rivaroxaban cohort.

In the secondary analysis of patients who adhered to the DOAC therapy interruption and resumption protocols, the 30-day postoperative rate of major bleeding was 1.2% (95% CI, 0%-1.89%) in the apixaban cohort, 1.0% (95% CI, 0%-1.93%) in the dabigatran cohort, and 1.69% (95% CI, 0%-2.53%) in the rivaroxaban cohort. The rate of arterial thromboembolism was 0.19% (95% CI, 0%-0.56%) in the apixaban cohort, 0.50% (95% CI, 0%-1.25%) in the dabigatran cohort, and 0.42% (95% CI, 0%-0.94%) in the rivaroxaban cohort (eAppendix 6 in the Supplement). Results according to clinical site are shown in eAppendix 7 in the Supplement.

Preoperative DOAC treatment levels were measured for 2541 patients (84.5%) (eAppendix 10 in the Supplement). The proportion of patients with a level less than 50 ng/mL was 90.5% in the apixaban cohort, 95.1% in the dabigatran cohort, and 96.8% in the rivaroxaban cohort. Among 1007 patients who had a high-bleeding-risk procedure, 832 (82.6%) had anticoagulant measurements, of whom the proportion with a residual anticoagulant level less than 50 ng/mL was 98.8%. The proportion of patients with a residual anticoagulant level of 30 to 49.9 ng/mL in the high-bleeding-risk procedure group was 4.8% in the apixaban cohort, 0.55% in the dabigatran cohort, and 14.0% in the rivaroxaban cohort. Results for the nonspecific coagulation tests are shown in eAppendix 8 in the Supplement.

Discussion

We found that in patients with AF who were receiving a DOAC (apixaban, dabigatran, or rivaroxaban) and required interruption of the anticoagulant regimen for elective surgery or procedure, a simple standardized perioperative management strategy that did not require the use of heparin bridging or preoperative coagulation function testing was associated with low rates of perioperative major bleeding (<2%) and arterial thromboembolism (<1%). Furthermore, a high proportion of patients (>90% overall; 98.8% of those at high bleeding risk) had a minimal or no residual anticoagulant level at the time of the procedure.

Based on the primary analysis cohort, our hypothesis that the PAUSE perioperative management strategy would exclude a 2% rate of major bleeding was supported in the dabigatran cohort (0.90%; 95% CI, 0%-1.73%) but not in the apixaban cohort (1.35%; 95% CI, 0%-2.0%) or rivaroxaban cohort (1.85%; 95% CI, 0%-2.65%), whereas our hypothesis that this management strategy would exclude a

Table 3. Primary Study Outcomes

Outcome	DOAC Cohort		
	Apixaban (n = 1257)	Dabigatran Etexilate (n = 668)	Rivaroxaban (n = 1082)
Primary			
Major bleeding ^a			
No. (%)	17 (1.35)	6 (0.90)	20 (1.85)
1-Sided 95% CI	0-2.00	0-1.73	0-2.65
P value	.051	.02	.36
Arterial thromboembolism ^{b,c}			
No. (%)	2 (0.16)	4 (0.60)	4 (0.37)
1-Sided 95% CI	0-0.48	0-1.33	0-0.82
P value	<.001	.03	.001
Secondary			
Death			
No. (%)	3 (0.24)	3 (0.45)	3 (0.28)
2-Sided 95% CI	0.08-0.70	0.15-1.31	0.09-0.81
Myocardial infarction			
No. (%)	1 (0.08)	0 (0)	0 (0)
2-Sided 95% CI	0.01-0.45	0-0.57	0-0.35
Deep vein thrombosis			
No. (%)	2 (0.16)	1 (0.15)	0 (0)
2-Sided 95% CI	0.04-0.58	0.03-0.84	0-0.35
Pulmonary embolism			
No. (%)	4 (0.32)	1 (0.15)	1 (0.09)
2-Sided 95% CI	0.12-0.82	0.03-0.84	0.02-0.52
Arterial catheter thrombosis ^d			
No. (%)	1 (0.08)	1 (0.15)	0 (0)
2-Sided 95% CI	0.01-0.45	0.03-0.84	0-0.35
Clinically relevant nonmajor bleeding			
No. (%)	21 (1.67)	13 (1.95)	26 (2.4)
2-Sided 95% CI	1.10-2.54	1.14-3.30	1.65-3.50
Minor bleeding			
No. (%)	54 (4.3)	38 (5.69)	62 (5.73)
2-Sided 95% CI	3.31-5.56	4.17-7.71	4.5-7.28

Abbreviation: DOAC, direct oral anticoagulant.

^a P value of the 1-sided test for 1 proportion to check that the proportion of major bleeding per DOAC was less than 2%.

^b P value of the 1-sided test for 1 proportion to check that the proportion of arterial thromboembolism per DOAC was less than 1.5%.

^c All episodes of arterial thromboembolism were ischemic stroke.

^d No episodes of catheter-related venous thrombosis were reported.

Table 4. Incidence of Major Bleeding by Elective Surgery or Procedure-Associated Bleeding Risk

Procedure-Associated Bleeding Risk	Apixaban Cohort (n = 1257)	Dabigatran Etexilate Cohort (n = 668)	Rivaroxaban Cohort (n = 1082)
Low bleeding risk			
No. (%)	851 (67.7)	440 (65.9)	709 (65.5)
30-d Postoperative rate of major bleeding, % (95% CI)	0.59 (0-1.20)	0.91 (0-2.01)	1.27 (0-2.17)
High bleeding risk			
No. (%)	406 (32.3)	228 (34.1)	373 (34.5)
30-d Postoperative rate of major bleeding, % (95% CI)	2.96 (0-4.68)	0.88 (0-2.62)	2.95 (0-4.76)

1.5% rate of arterial thromboembolism was supported in all 3 cohorts. In the per protocol analysis, excluding a 2% rate of major bleeding was supported in the dabigatran cohort (1.0%; 95% CI, 0%-1.93%) and the apixaban cohort (1.2%; 95% CI, 0%-1.89%) but not in the rivaroxaban cohort (1.69%; 95% CI, 0%-2.53%); excluding a rate of arterial thromboembolism of 1.5% was supported in all 3 cohorts.

Our exploratory postulation that a high proportion of patients (>90%) would have a residual anticoagulant level less

than 50 ng/mL at the time of the operation was supported in all 3 DOAC cohorts. In addition, we found that, among patients with a high-bleeding-risk procedure (which included any patient with neuraxial anesthesia) in whom there was concern of bleeding complications associated with an excessive residual anticoagulant level,^{16,18,19} almost all patients (98.8%) had a residual anticoagulant level less than 50 ng/mL. Moreover, the proportion of such patients with a residual anticoagulant level less than 30 mg/mL, which some experts con-

sider an optimal preoperative anticoagulant level,¹³ was high in the apixaban cohort (93.1%) and dabigatran cohort (98.9%). Among patients in the rivaroxaban cohort with a high-bleeding-risk procedure, a lower proportion (85.4%) had a residual anticoagulant level less than 30 ng/mL, an observation that requires further study. When the findings were assessed according to procedure-associated high- and low-bleeding risk, rates of major bleeding appeared to be higher among patients with a high-bleeding-risk procedure in the apixaban and rivaroxaban cohorts. This finding may reflect an intrinsically higher rate of bleeding expected with major procedures. Further study is needed to assess the PAUSE perioperative DOAC regimen management in patients with high-bleeding-risk procedures.

Most other studies that assessed perioperative DOAC regimen management are not comparable to this study because their management was not standardized, perioperative heparin bridging was allowed, and fewer patients (10%-20%) with high bleeding risk were studied.^{3,10-14} In these studies, perioperative rates of major bleeding, for example, varied and were as high as 6%.³ Two pertinent studies that assessed standardized perioperative anticoagulant management without heparin bridging had similar adverse outcome rates to those in this study. In a cohort study of 541 patients receiving dabigatran who had standardized perioperative interruption and resumption of their treatment, the 30-day postoperative rate of major bleeding was 1.8% and the arterial thromboembolism rate was 0.2%.¹⁴ In the BRIDGE trial, which evaluated a bridging strategy in patients with AF who had perioperative warfarin treatment interruption, patients who were not bridged had a 30-day postoperative rate of major bleeding of 1.3% and arterial thromboembolism rate of 0.4%,⁷ and those who underwent a high-bleeding-risk procedure had a rate of major bleeding of 3.2%.³⁴

Limitations and Strengths

This study has limitations. First, although a cohort study design may introduce patient selection bias, this was unlikely because a high proportion (83%) of screened patients participated in this study, and their risk factor profile, as measured by the CHA₂DS₂-VASc risk score, was comparable to that of patients with AF included in population-based studies.³⁵ Second, although few patients (n = 230) received neuraxial anesthesia, in whom there was a concern about bleeding risk during

the operation associated with an excessive residual anticoagulant level, the management of such patients was the same as all patients undergoing a high-bleeding-risk procedure (n = 1007). Accordingly, the high proportion (98.8%) of patients with minimal to no residual anticoagulant level in this group would be applicable to those with neuraxial anesthesia. Third, the dabigatran cohort did not reach the expected sample size, owing to the decrease in dabigatran use compared with other DOAC regimens during the study, but the number of patients accrued was sufficient to address the study hypotheses in this cohort. Fourth, patients using edoxaban tosylate were not included as the drug was not available for clinical use when the PAUSE study started, and the results are not generalizable to this DOAC. Fifth, the 50 ng/mL cut point used in this study to define a clinically important residual preoperative DOAC level was not established, and further study is needed to assess a correlation between preoperative DOAC treatment levels and bleeding. Sixth, most patients included were white, and additional studies are needed in nonwhite populations. Seventh, patients with venous thromboembolism, who represent a different study population,³⁶ were not included.

A strength of this study is the generalizability of the results to patients assessed in clinical practice, as a high proportion of screened patients were enrolled (83%) and few were lost to follow-up (<1%). Another strength is the clinical applicability of the DOAC regimen management we assessed, as most patients adhered to the perioperative DOAC therapy interruption (95%) and resumption (93%) management protocol. The simple strategy of omitting DOAC regimens for 1 day before and after a low-bleeding-risk procedure and 2 days before and after a high-bleeding-risk procedure (except for patients using dabigatran with a CrCl <50 mL/min) is, therefore, likely to be easily adoptable in clinical practice.

Conclusions

In this study, patients with AF who had DOAC therapy interruption for elective surgery or procedure, a simple standardized perioperative management strategy without heparin bridging or measurement of coagulation function was associated with low rates of major bleeding and arterial thromboembolism.

ARTICLE INFORMATION

Accepted for Publication: May 13, 2019.

Published Online: August 5, 2019.

doi:10.1001/jamainternmed.2019.2431

Author Affiliations: Department of Medicine, McMaster University, Hamilton, Ontario, Canada (Douketis, Duncan, Gross, Bates, Spencer, Arnold, Li, Schulman); The Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, Department of Medicine, Northwell Health at Lenox Hill Hospital, New York, New York (Spyropoulos); Ottawa Hospital Research Institute, Department of Medicine, University of Ottawa, Ottawa, Ontario, Canada (Carrier); L'Institut

du Savoir Montfort, L'Hopital Montfort, Ottawa, Ontario, Canada (Le Gal); Department of Surgery, NorthShore University Health Systems, Evanston, Illinois (Tafur, Caprini); Department of Cardiovascular Sciences, University of Leuven, Leuven, Belgium (Vanassche, Verhamme); Department of Medicine, Dalhousie University, Halifax, Nova Scotia, Canada (Shivakumar); Department of Medicine, University of British Columbia, Vancouver, British Columbia, Canada (Lee); Department of Medicine, University of Toronto, Toronto, Ontario, Canada (Yeo); Department of Medicine, Montreal General Hospital, McGill University, Montreal, Quebec, Canada (Solymoss); Department of Medicine,

Université de Montréal, Montreal, Quebec, Canada (Kassis); Department of Internal Medicine, Centre Hospitalier Universitaire de Sherbrooke, Sherbrooke, Quebec, Canada (Le Templier); Department of Anesthesiology, University of Manitoba, Winnipeg, Manitoba, Canada (Kowalski); Department of Medicine, Jewish General Hospital, McGill University, Montreal, Quebec, Canada (Blostein); Department of Medicine, Henry Ford Hospital, Detroit, Michigan (Shah); Department of Medicine, University of Calgary, Calgary, Alberta, Canada (MacKay); Department of Medicine, University of Alberta, Edmonton, Alberta, Canada (Wu); Department of Pharmacy, Kaiser Permanente Colorado, Aurora, Colorado (Clark); Department of

Anesthesiology, University of Thessaly, Larissa, Greece (Arnaoutoglou); Department of Vascular Medicine, Amsterdam Cardiovascular Sciences, Amsterdam UMC, Location AMC, the Netherlands (Coppens); Hamilton Regional Laboratory Medicine Program, McMaster University, Hamilton, Ontario, Canada (Moffat); Department of Anesthesiology, Hamilton Health Sciences, McMaster University, Hamilton, Ontario, Canada (Syed); Department of Obstetrics and Gynecology, The First I.M. Sechenov Moscow State Medical University, Moscow, Russia (Schulman).

Author Contributions: Drs Spyropoulos and Schulman had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. The PAUSE steering committee comprised 5 standing members, Drs Douketis, Caprini, Schulman, Spyropoulos, and Syed, and 3 to 4 rotating site investigators.

Concept and design: Douketis, Spyropoulos, Verhamme, Lee, Kassis, Shah, Wu, Syed, Schulman.

Acquisition, analysis, or interpretation of data: Douketis, Spyropoulos, Duncan, Carrier, Le Gal, Tafur, Vanassche, Shivakumar, Gross, Lee, Yeo, Solymoss, Le Templier, Kowalski, Blostein, MacKay, Clark, Bates, Spencer, Arnaoutoglou, Coppens, Arnold, Caprini, Li, Moffat, Syed.

Drafting of the manuscript: Douketis, Spyropoulos, Li. **Critical revision of the manuscript for important intellectual content:** All authors.

Statistical analysis: Douketis, Li.

Obtained funding: Douketis, Carrier, Gross, Lee, MacKay, Schulman.

Administrative, technical, or material support: Douketis, Duncan, Carrier, Le Gal, Vanassche, Gross, Kassis, Kowalski, Clark, Bates, Arnold, Moffat, Syed. **Supervision:** Douketis, Spyropoulos, Duncan, Tafur, Verhamme, Gross, Yeo, Blostein, Shah, MacKay, Caprini, Moffat.

Other - Principal investigator: Douketis.

Other - major part of the patient recruitment: Schulman.

Other - contribution in data collection: Arnaoutoglou.

Other - Laboratory support: Moffat.

Other - member of steering committee: Syed.

Conflict of Interest Disclosures: Dr Douketis reported personal fees from Pfizer, Sanofi, Leo Pharma, Bristol-Myers Squibb, Janssen, The Merck Manual, and UpToDate outside of the submitted work. Dr Spyropoulos reported grants and personal fees from Janssen and Boehringer Ingelheim and personal fees from Bayer, Portola, and ATLAS Group outside of the submitted work. Dr Carrier reported grants from Canadian Institutes of Health Research (CIHR) during the conduct of the study; grants from Leo Pharma and BMS; and personal fees from Pfizer; and personal fees from Bayer, Servier, BMS, and Leo Pharma outside of the submitted work. Dr Le Gal reported personal fees from Bayer, Pfizer, Leo Pharma, Sanofi, and bioMérieux as well as other fees from Portola Pharmaceuticals, Boehringer-Ingelheim, Pfizer, Bristol-Myers Squibb, Leo Pharma, Daiichi Sankyo, and Bayer outside of the submitted work. Dr Tafur reported grants from PAUSE during the conduct of the study. Dr Vanassche reported other fees from Bayer, Boehringer Ingelheim, Daiichi Sankyo, and Leo Pharma outside of the submitted work. Dr Verhamme reported grants and personal fees from Bayer HealthCare, Boehringer Ingelheim, Pfizer, BMS, Daiichi Sankyo, and Leo Pharma as well

as personal fees from Portola outside of the submitted work. Dr Shivakumar reported personal fees from Pfizer Inc and Bayer Inc outside of the submitted work. Dr Gross reported other fees from Bayer and Pfizer; grants and other fees from Bristol-Myers-Squibb; and grants from Boehringer-Ingelheim during the conduct of the study; other fees from Servier, Leo Pharrma, and Alexion outside of the submitted work; and a patent to US10131932B2 method for assaying a protease issued. Dr Lee reported personal fees from Bayer, Leo Pharma, and Pfizer as well as grants, personal fees, and nonfinancial support from BMS during the conduct of the study. Dr Le Templier reported personal fees from BMS-Pfizer outside of the submitted work. Dr Shah reported grants from CIHR during the conduct of the study. Dr MacKay reported grants from CIHR during the conduct of the study and personal fees from BMS/Pfizer outside of the submitted work. Dr Wu reported other fees from Leo Pharma, Pfizer, Servier, BMS-Pfizer, Bayer, and Daiichi-Sankyo during the conduct of the study. Dr Bates reported grant 313156 from CIHR and grant G-14-0006136 from Heart and Stroke Foundation of Canada during the conduct of the study as well as grants from Bayer Inc outside of the submitted work. Dr Arnaoutoglou reported grants and personal fees from Bayer outside of the submitted work. Dr Coppens reported other fees from Bristol-Myers Squibb, Pfizer, and Boehringer-Ingelheim; grants, personal fees, and other fees from Bayer and Daiichi Sankyo; personal fees from Portola; and grants from Sanquin Blood Supply outside of the submitted work. Dr Caprini reported personal fees from BMS, Janssen, and Pfizer outside of the submitted work. Dr Syed reported personal fees and other fees from Octapharma outside of the submitted work. Dr Schulman reported grants and personal fees from Boehringer Ingelheim, Octapharma as well as personal fees from Bayer, Daiichi Sankyo, Pfizer, Alnylam, and Sanofi during the conduct of the study. No other disclosures were reported.

Funding/Support: The PAUSE study was funded by grant 313156 from the Canadian Institutes of Health Research and grant G-14-0006136 from the Heart and Stroke Foundation of Canada. Additional support was provided by the CanVECTOR research network. In-kind support was obtained from Aniera-Hyphen Biomed, which provided the anti-factor Xa and dilute thrombin time assays.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: We thank the members of the Data Safety Monitoring Board: Walter Ageno, MD; David Garcia, MD (chair), and Lehana Thabane, PhD. We also thank the members of the Event Adjudication Committee: Wendy Lim, MD; Lori Linkins, MD; William Risteovski, MD; and Demetrios J. Sahlas, MD. We are grateful to the research assistants from the McMaster Centre for Transfusion Research, Kayla Lucier, Grace Wang, and Tara McDougall, as well as the members of the McMaster University Clinical Research Laboratory and Biobank and Special Coagulation Laboratory at McMaster University Medical Center. We extend our gratitude to the clinical center study

coordinators and the patients who participated in this study. These individuals received no additional compensation, outside of their usual salary, for their contributions.

REFERENCES

1. Spyropoulos AC, Al-Badri A, Sherwood MW, Douketis JD. Perioperative management of patients receiving a vitamin K antagonist or a direct oral anticoagulant requiring an elective procedure or surgery. *J Thromb Haemost.* 2016;14(5):875-885. doi:10.1111/jth.13305
2. Zulkifly H, Lip GH, Lane DA. Epidemiology of atrial fibrillation. *Int J Clin Pract.* 2018;72(3):e13070. doi:10.1111/ijcp.13070
3. Healey JS, Eikelboom J, Douketis J, et al; RE-LY Investigators. Perioperative bleeding and thromboembolic events with dabigatran compared with warfarin: results from the Randomized Evaluation of Long-Term Anticoagulation Therapy (RE-LY) randomized trial [published correction appears in *Circulation.* 2012;126(10):e160]. *Circulation.* 2012;126(3):343-348. doi:10.1161/CIRCULATIONAHA.111.090464
4. Spyropoulos AC, Douketis JD. How I treat anticoagulated patients undergoing an elective procedure or surgery. *Blood.* 2012;120(15):2954-2962. doi:10.1182/blood-2012-06-415943
5. Dunn AS, Spyropoulos AC, Turpie AG. Bridging therapy in patients on long-term oral anticoagulants who require surgery: the Prospective Peri-operative Enoxaparin Cohort Trial (PROSPECT). *J Thromb Haemost.* 2007;5(11):2211-2218. doi:10.1111/j.1538-7836.2007.02729.x
6. Douketis JD, Healey JS, Brueckmann M, et al. Perioperative bridging anticoagulation during dabigatran or warfarin interruption among patients who had an elective surgery or procedure. Substudy of the RE-LY trial. *Thromb Haemost.* 2015; 113(3):625-632. doi:10.1160/TH14-04-0305
7. Douketis JD, Spyropoulos AC, Kaatz S, et al; BRIDGE Investigators. Perioperative bridging anticoagulation in patients with atrial fibrillation. *N Engl J Med.* 2015;373(9):823-833. doi:10.1056/NEJMoa1501035
8. Tripodi A. To measure or not to measure direct oral anticoagulants before surgery or invasive procedures: reply. *J Thromb Haemost.* 2016;14(12):2559-2561. doi:10.1111/jth.13513
9. Spyropoulos AC, Al-Badri A, Sherwood MW, Douketis JD. To measure or not to measure direct oral anticoagulants before surgery or invasive procedures: comment. *J Thromb Haemost.* 2016;14(12):2556-2559. doi:10.1111/jth.13505
10. Garcia D, Alexander JH, Wallentin L, et al. Management and clinical outcomes in patients treated with apixaban vs warfarin undergoing procedures. *Blood.* 2014;124(25):3692-3698. doi:10.1182/blood-2014-08-595496
11. Sherwood MW, Douketis JD, Patel MR, et al; ROCKET AF Investigators. Outcomes of temporary interruption of rivaroxaban compared with warfarin in patients with nonvalvular atrial fibrillation: results from the rivaroxaban once daily, oral, direct factor Xa inhibition compared with vitamin K antagonism for prevention of stroke and embolism trial in atrial fibrillation (ROCKET AF). *Circulation.* 2014;129(18):1850-1859. doi:10.1161/CIRCULATIONAHA.113.005754

12. Beyer-Westendorf J, Gelbricht V, Förster K, et al. Peri-interventional management of novel oral anticoagulants in daily care: results from the prospective Dresden NOAC registry. *Eur Heart J*. 2014;35(28):1888-1896. doi:10.1093/eurheartj/ehx557
13. Godier A, Dincq AS, Martin AC, et al. Predictors of pre-procedural concentrations of direct oral anticoagulants: a prospective multicentre study. *Eur Heart J*. 2017;38(31):2431-2439. doi:10.1093/eurheartj/ehx403
14. Schulman S, Carrier M, Lee AY, et al; Periop Dabigatran Study Group. Perioperative management of dabigatran: a prospective cohort study. *Circulation*. 2015;132(3):167-173. doi:10.1161/CIRCULATIONAHA.115.015688
15. Faraoni D, Samama CM, Ranucci M, Dietrich W, Levy JH. Perioperative management of patients receiving new oral anticoagulants: an international survey. *Clin Lab Med*. 2014;34(3):637-654. doi:10.1016/j.cl.2014.06.006
16. Narouze S, Benzon HT, Provenzano DA, et al. Interventional spine and pain procedures in patients on antiplatelet and anticoagulant medications: guidelines from the American Society of Regional Anesthesia and Pain Medicine, the European Society of Regional Anaesthesia and Pain Therapy, the American Academy of Pain Medicine, the International Neuromodulation Society, the North American Neuromodulation Society, and the World Institute of Pain. *Reg Anesth Pain Med*. 2015;40(3):182-212. doi:10.1097/AAP.0000000000000223
17. Raval AN, Cigarroa JE, Chung MK, et al; American Heart Association Clinical Pharmacology Subcommittee of the Acute Cardiac Care and General Cardiology Committee of the Council on Clinical Cardiology; Council on Cardiovascular Disease in the Young; and Council on Quality of Care and Outcomes Research. Management of patients on non-vitamin K antagonist oral anticoagulants in the acute care and periprocedural setting: a scientific statement from the American Heart Association. *Circulation*. 2017;135(10):e604-e633. doi:10.1161/CIR.0000000000000477
18. Albaladejo P, Bonhomme F, Blais N, et al; French Working Group on Perioperative Hemostasis (GIHP). Management of direct oral anticoagulants in patients undergoing elective surgeries and invasive procedures: updated guidelines from the French Working Group on Perioperative Hemostasis (GIHP) - September 2015. *Anaesth Crit Care Pain Med*. 2017;36(1):73-76. doi:10.1016/j.jccpm.2016.09.002
19. Horlocker TT, Vandermeulen E, Kopp SL, Gogarten W, Leffert LR, Benzon HT. Regional Anesthesia in the Patient Receiving Antithrombotic or Thrombolytic Therapy: American Society of Regional Anesthesia and Pain Medicine Evidence-Based Guidelines (Fourth Edition). *Reg Anesth Pain Med*. 2018;43(3):263-309. doi:10.1097/AAP.0000000000000763
20. Steffel J, Verhamme P, Potpara TS, et al; ESC Scientific Document Group. The 2018 European Heart Rhythm Association practical guide on the use of non-vitamin K antagonist oral anticoagulants in patients with atrial fibrillation. *Eur Heart J*. 2018;39(16):1330-1393. doi:10.1093/eurheartj/ehy136
21. Siegal D, Yudin J, Kaatz S, Douketis JD, Lim W, Spyropoulos AC. Periprocedural heparin bridging in patients receiving vitamin K antagonists: systematic review and meta-analysis of bleeding and thromboembolic rates. *Circulation*. 2012;126(13):1630-1639. doi:10.1161/CIRCULATIONAHA.112.105221
22. Douketis JD, Spyropoulos AC, Anderson JM, et al. The perioperative anticoagulant use for surgery evaluation (PAUSE) study for patients on a direct oral anticoagulant who need an elective surgery or procedure: design and rationale. *Thromb Haemost*. 2017;117(12):2415-2424. doi:10.1160/TH17-08-0553
23. Vandembroucke JP, von Elm E, Altman DG, et al; STROBE Initiative. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Epidemiology*. 2007;18(6):805-835. doi:10.1097/EDE.0b013e3181577511
24. Euser AM, Zoccali C, Jager KJ, Dekker FW. Cohort studies: prospective versus retrospective. *Nephron Clin Pract*. 2009;113(3):c214-c217. doi:10.1159/000235241
25. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron*. 1976;16(1):31-41. doi:10.1159/000180580
26. Doherty JU, Gluckman TJ, Hucker WJ, et al. 2017 ACC expert consensus decision pathway for periprocedural management of anticoagulation in patients with nonvalvular atrial fibrillation: a report of the American College of Cardiology Clinical Expert Consensus Document Task Force. *J Am Coll Cardiol*. 2017;69(7):871-898. doi:10.1016/j.jacc.2016.11.024
27. Douketis JD, Spyropoulos AC, Spencer FA, et al. Perioperative management of antithrombotic therapy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. [published correction appears in *Chest*. 2012;141(4):1129]. *Chest*. 2012;141(2)(suppl):e326S-e350S. doi:10.1378/chest.11-2298
28. Schulman S, Angerås U, Bergqvist D, Eriksson B, Lassen MR, Fisher W; Subcommittee on Control of Anticoagulation of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in surgical patients. *J Thromb Haemost*. 2010;8(1):202-204. doi:10.1111/j.1538-7836.2009.03678.x
29. Spyropoulos AC, Douketis JD, Gerotziafas G, Kaatz S, Ortel TL, Schulman S; Subcommittee on Control of Anticoagulation of the SSC of the ISTH. Periprocedural antithrombotic and bridging therapy: recommendations for standardized reporting in patients with arterial indications for chronic oral anticoagulant therapy. *J Thromb Haemost*. 2012;10(4):692-694. doi:10.1111/j.1538-7836.2012.04630.x
30. Samuelson BT, Cuker A, Siegal DM, Crowther M, Garcia DA. Laboratory assessment of the anticoagulant activity of direct oral anticoagulants: a systematic review. *Chest*. 2017;151(1):127-138. doi:10.1016/j.chest.2016.08.1462
31. Tripodi A, Padovan L, Veena C, Scalabrino E, Testa S, Peyvandi F. How the direct oral anticoagulant apixaban affects thrombin generation parameters. *Thromb Res*. 2015;135(6):1186-1190. doi:10.1016/j.thromres.2015.03.032
32. Dunn AS, Turpie AG. Perioperative management of patients receiving oral anticoagulants: a systematic review. *Arch Intern Med*. 2003;163(8):901-908. doi:10.1001/archinte.163.8.901
33. Wallis S. Binomial confidence intervals and contingency tests: mathematical fundamentals and the evaluation of alternative methods. *J Quant Linguist*. 2013;20(3):178-208. doi:10.1080/09296174.2013.799918
34. Clark NP, Douketis JD, Hasselblad V, Schulman S, Kindzelski AL, Ortel TL; BRIDGE Investigators. Predictors of perioperative major bleeding in patients who interrupt warfarin for an elective surgery or procedure: analysis of the BRIDGE trial. *Am Heart J*. 2018;195:108-114. doi:10.1016/j.ahj.2017.09.015
35. Forslund T, Komen JJ, Andersen M, et al. Improved stroke prevention in atrial fibrillation after the introduction of non-vitamin K antagonist oral anticoagulants. *Stroke*. 2018;49(9):2122-2128. doi:10.1161/STROKEAHA.118.021990
36. Clark NP, Witt DM, Davies LE, et al. Bleeding, recurrent venous thromboembolism, and mortality risks during warfarin interruption for invasive procedures. *JAMA Intern Med*. 2015;175(7):1163-1168. doi:10.1001/jamainternmed.2015.1843