

CLINICAL PRACTICE GUIDELINE: EXECUTIVE SUMMARY

2021 AHA/ACC/ASE/CHEST/SAEM/SCCT/SCMR Guideline for the Evaluation and Diagnosis of Chest Pain: Executive Summary



A Report of the American College of Cardiology/American Heart Association
Joint Committee on Clinical Practice Guidelines

**Writing
Committee
Members***

Martha Gulati, MD, MS, FACC, FAHA, *Chair*†
Phillip D. Levy, MD, MPH, FACC, FAHA, *Vice Chair*†
Debabrata Mukherjee, MD, MS, FACC, FAHA, *Vice Chair*†
Ezra Amsterdam, MD, FACC†
Deepak L. Bhatt, MD, MPH, FACC, FAHA†
Kim K. Birtcher, MS, PHARM, AACCF†
Ron Blankstein, MD, FACC, MSCCT§
Jack Boyd, MD†
Renee P. Bullock-Palmer, MD, FACC, FAHA, FASE, FSCCT†
Theresa Conejo, RN, BSN, FAHA||
Deborah B. Diercks, MD, MSc, FACC¶
Federico Gentile, MD, FACC#
John P. Greenwood, MBChB, PhD, FSCMR, FACC**
Erik P. Hess, MD, MSc†
Steven M. Hollenberg, MD, FACC, FAHA, FCCP††
Wael A. Jaber, MD, FACC, FASE‡‡

Hani Jneid, MD, FACC, FAHA§§
José A. Joglar, MD, FAHA, FACC‡
David A. Morrow, MD, MPH, FACC, FAHA†
Robert E. O'Connor, MD, MPH, FAHA†
Michael A. Ross, MD, FACC†
Leslee J. Shaw, PhD, FACC, FAHA, MSCCT†

*Writing committee members are required to recuse themselves from voting on sections to which their specific relationships with industry may apply; see [Appendix 1](#) for detailed information. †ACC/AHA Representative. ‡ACC/AHA Joint Committee on Clinical Practice Guidelines Liaison. §Society of Cardiovascular Computed Tomography Representative. ||Lay Patient Representative. ¶Society for Academic Emergency Medicine Representative. #Former ACC/AHA Joint Committee member; current member during the writing effort. **Society for Cardiovascular Magnetic Resonance Representative. ††American College of Chest Physicians Representative. ‡‡American Society of Echocardiography Representative. §§Task Force on Performance Measures, Liaison.

This document was approved by the American College of Cardiology Clinical Policy Approval Committee in May 2021, the American Heart Association Science Advisory and Coordinating Committee in May 2021, the Society of Cardiovascular Computed Tomography in July 2021, the Society for Academic Emergency Medicine Representative in June 2021, the Society for Cardiovascular Magnetic Resonance in June 2021, the American College of Chest Physicians in June 2021, the American Society of Echocardiography in June 2021, the American Heart Association Executive Committee in July 2021, and the American College of Cardiology Science and Quality Committee in July 2021.

The American College of Cardiology requests that this document be cited as follows: Gulati M, Levy PD, Mukherjee D, Amsterdam E, Bhatt DL, Birtcher KK, Blankstein R, Boyd J, Bullock-Palmer RP, Conejo T, Diercks DB, Gentile F, Greenwood JP, Hess EP, Hollenberg SM, Jaber WA, Jneid H, Joglar JA, Morrow DA, O'Connor RE, Ross MA, Shaw LJ. 2021 AHA/ACC/ASE/CHEST/SAEM/SCCT/SCMR guideline for the evaluation and diagnosis of chest pain: executive summary: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2021;78:(22):2218-2261.

This article has been copublished in *Circulation*.

Copies: This document is available on the websites of the American College of Cardiology (www.acc.org) and the American Heart Association (<https://professional.heart.org/>). For copies of this document, please contact the Elsevier Inc. Reprint Department via fax (212-633-3820) or e-mail (reprints@elsevier.com).

Permissions: Multiple copies, modification, alteration, enhancement, and/or distribution of this document are not permitted without the express permission of the American College of Cardiology. Requests may be completed online via the Elsevier site (<http://www.elsevier.com/about/policies/author-agreement/obtaining-permission>).

ACC/AHA Joint Committee Members

Patrick T. O’Gara, MD, MACC, FAHA, *Chair*
 Joshua A. Beckman, MD, MS, FAHA, FACC,
Chair-Elect
 Glenn N. Levine, MD, FACC, FAHA,
Immediate Past Chair#

Sana M. Al-Khatib, MD, MHS, FACC, FAHA#
 Anastasia L. Armbruster, PharmD, FACC
 Kim K. Birtcher, MS, PharmD, AACC#
 Ralph G. Brindis, MD, MPH, MACC#
 Joaquin E. Cigarroa, MD, FACC#
 Lisa de las Fuentes, MD, MS, FASE, FAHA
 Anita Deswal, MD, MPH, FACC, FAHA
 Dave L. Dixon, PharmD, FACC#
 Lee A. Fleisher, MD, FACC, FAHA#
 Federico Gentile, MD, FACC#
 Zachary D. Goldberger, MD, MS, FACC, FAHA#
 Bulent Gorenek, MD, FACC
 Norrisa Haynes, MD

Adrian F. Hernandez, MD
 Mark A. Hlatky, MD, FACC, FAHA#
 John S. Ikonomidis, MD, PhD, FAHA#
 José A. Joglar, MD, FAHA, FACC
 W. Schuyler Jones, MD, FACC
 Joseph E. Marine, MD, FACC#
 Daniel B. Mark, MD, MPH, FACC
 Debabrata Mukherjee, MD, MS, FACC, FAHA
 Latha P. Palaniappan, MD, MS, FAHA, FACC
 Mariann R. Piano, RN, PhD, FAHA
 Tanveer Rab, MD, FACC
 Barbara Riegel, PhD, RN, FAHA#
 Erica S. Spatz, MD, MHS, FACC
 Jacqueline E. Tamis-Holland, MD, FACC
 Duminda N. Wijeyesundera, MD, PhD#
 Y. Joseph Woo, MD, FAHA, FACC

#Former ACC/AHA Joint Committee member; current member during the writing effort.

ABSTRACT

AIM This executive summary of the clinical practice guideline for the evaluation and diagnosis of chest pain provides recommendations and algorithms for clinicians to assess and diagnose chest pain in adult patients.

METHODS A comprehensive literature search was conducted from November 11, 2017, to May 1, 2020, encompassing studies, reviews, and other evidence conducted on human subjects that were published in English from PubMed, EMBASE, the Cochrane Collaboration, Agency for Healthcare Research and Quality reports, and other relevant databases. Additional relevant studies, published through April 2021, were also considered.

STRUCTURE Chest pain is a frequent cause for emergency department visits in the United States. The “2021 AHA/ACC/AASE/CHEST/SAEM/SCCT/SCMR Guideline for the Evaluation and Diagnosis of Chest Pain” provides recommendations based on contemporary evidence on the assessment and evaluation of chest pain. These guidelines present an evidence-based approach to risk stratification and the diagnostic workup for the evaluation of chest pain. Cost-value considerations in diagnostic testing have been incorporated and shared decision-making with patients is recommended.

CONTENTS

ABSTRACT	2219	2.1. History	2223
TOP 10 TAKE-HOME MESSAGES FOR THE EVALUATION AND DIAGNOSIS OF CHEST PAIN ...	2220	2.1.1. A Focus on the Uniqueness of Chest Pain in Women	2225
1. PURPOSE OF THE EXECUTIVE SUMMARY	2221	2.1.2. Considerations for Older Patients With Chest Pain	2225
1.1. Document Review and Approval	2222	2.1.3. Considerations for Diverse Patient Populations With Chest Pain	2225
1.2. Class of Recommendations and Level of Evidence	2222	2.1.4. Patient-Centric Considerations	2225
1.3. Defining Chest Pain	2223	2.2. Physical Examination	2226
2. INITIAL EVALUATION	2223	2.3. Diagnostic Testing	2226
		2.3.1. Setting Considerations	2226
		2.3.2. Electrocardiogram	2227

2.3.3. Chest Radiography	2228
2.3.4. Biomarkers	2228
3. CARDIAC TESTING GENERAL CONSIDERATIONS	2229
4. CHOOSING THE RIGHT PATHWAY WITH PATIENT-CENTRIC ALGORITHMS FOR ACUTE CHEST PAIN	2232
4.1. Patients With Acute Chest Pain and Suspected Acute Coronary Syndrome (Not Including STEMI)	2233
4.1.1. Low-Risk Patients With Acute Chest Pain	2236
4.1.2. Intermediate-Risk Patients With Acute Chest Pain	2237
4.1.3. High-Risk Patients With Acute Chest Pain	2240
4.1.4. Acute Chest Pain in Patients With Prior Coronary Artery Bypass Graft (CABG) Surgery	2240
4.1.5. Evaluation of Patients With Acute Chest Pain Receiving Dialysis	2240
4.1.6. Evaluation of Acute Chest Pain in Patients With Cocaine and Methamphetamine Use	2240
4.1.7. Shared Decision-Making in Patients With Acute Chest Pain	2241
4.2. Evaluation of Acute Chest Pain With Nonischemic Cardiac Pathologies	2241
4.2.1. Acute Chest Pain With Suspected Acute Aortic Syndrome	2241
4.2.2. Acute Chest Pain With Suspected PE	2241
4.2.3. Acute Chest Pain With Suspected Myopericarditis	2242
4.2.4. Acute Chest Pain With Valvular Heart Disease (VHD)	2242
4.3. Evaluation of Acute Chest Pain With Suspected Noncardiac Causes	2243
4.3.1. Evaluation of Acute Chest Pain With Suspected Gastrointestinal Syndromes	2244
4.3.2. Evaluation of Acute Chest Pain With Suspected Anxiety and Other Psychosomatic Considerations	2244
4.3.3. Evaluation of Acute Chest Pain in Patients With Sickle Cell Disease	2244
5. EVALUATION OF PATIENTS WITH STABLE CHEST PAIN	2244
5.1. Patients With No Known CAD Presenting With Stable Chest Pain	2244
5.1.2. Low-Risk Patients With Stable Chest Pain and No Known CAD	2245
5.1.3. Intermediate-High Risk Patients With Stable Chest Pain and No Known CAD	2246
5.2. Patients With Known CAD Presenting With Stable Chest Pain	2248

5.2.1. Patients With Obstructive CAD Who Present With Stable Chest Pain	2248
5.2.2. Patients With Known Nonobstructive CAD Presenting With Stable Chest Pain	2250
5.2.3. Patients With Suspected Ischemia and No Obstructive CAD (INOCA)	2251

REFERENCES	2252
-----------------------------	-------------

TOP 10 TAKE-HOME MESSAGES FOR THE EVALUATION AND DIAGNOSIS OF CHEST PAIN

- Chest Pain Means More Than Pain in the Chest.** Pain, pressure, tightness, or discomfort in the chest, shoulders, arms, neck, back, upper abdomen, or jaw, as well as shortness of breath and fatigue should all be considered anginal equivalents.
- High-Sensitivity Troponins Preferred.** High-sensitivity cardiac troponins are the preferred standard for establishing a biomarker diagnosis of acute myocardial infarction, allowing for more accurate detection and exclusion of myocardial injury.
- Early Care for Acute Symptoms.** Patients with acute chest pain or chest pain equivalent symptoms should seek medical care immediately by calling 9-1-1. Although most patients will not have a cardiac cause, the evaluation of all patients should focus on the early identification or exclusion of life-threatening causes.
- Share the Decision-Making.** Clinically stable patients presenting with chest pain should be included in decision-making; information about risk of adverse events, radiation exposure, costs, and alternative options should be provided to facilitate the discussion.
- Testing Not Needed Routinely for Low-Risk Patients.** For patients with acute or stable chest pain determined to be low risk, urgent diagnostic testing for suspected coronary artery disease is not needed.
- Pathways.** Clinical decision pathways for chest pain in the emergency department and outpatient settings should be used routinely.
- Accompanying Symptoms.** Chest pain is the dominant and most frequent symptom for both men and women ultimately diagnosed with acute coronary syndrome. Women may be more likely to present with accompanying symptoms such as nausea and shortness of breath.
- Identify Patients Most Likely to Benefit From Further Testing.** Patients with acute or stable chest pain who are at intermediate risk or intermediate to high pre-test risk of obstructive coronary artery disease, respectively, will benefit the most from cardiac imaging and testing.
- Noncardiac Is In. Atypical Is Out.** “Noncardiac” should be used if heart disease is not suspected.



“Atypical” is a misleading descriptor of chest pain, and its use is discouraged.

- 10. Structured Risk Assessment Should Be Used.** For patients presenting with acute or stable chest pain, risk for coronary artery disease and adverse events should be estimated using evidence-based diagnostic protocols.

Figure 1 illustrates the take-home messages.

1. PURPOSE OF THE EXECUTIVE SUMMARY

The charge of the writing committee was to develop a guideline for the evaluation of acute or stable chest pain or other anginal equivalents, in a variety of clinical settings, with an emphasis on the diagnosis on ischemic causes (1). The guideline will not provide

recommendations on whether revascularization is appropriate, or what modality is indicated (1). Such recommendations can be found in the forthcoming American Heart Association (AHA)/American College of Cardiology (ACC) coronary artery revascularization guideline (1a).

After injuries, chest pain is the second most common reason for adults to present to the emergency department (ED) in the United States and accounts for >6.5 million visits, which is 4.7% of all ED visits (2). Chest pain also leads to nearly 4 million outpatient visits annually in the United States (3). Chest pain remains a diagnostic challenge in the ED and outpatient setting and requires thorough clinical evaluation. Although the cause of chest pain is often noncardiac, coronary artery disease (CAD) affects >18.2 million adults in the United States and remains the leading cause of death for men and women,

TABLE 1 Applying ACC/AHA Class of Recommendation and Level of Evidence to Clinical Strategies, Interventions, Treatments, or Diagnostic Testing in Patient Care (Updated May 2019)*

CLASS (STRENGTH) OF RECOMMENDATION	LEVEL (QUALITY) OF EVIDENCE†
CLASS 1 (STRONG) Benefit >>> Risk Suggested phrases for writing recommendations: <ul style="list-style-type: none"> • Is recommended • Is indicated/useful/effective/beneficial • Should be performed/administered/other • Comparative-Effectiveness Phrases‡: <ul style="list-style-type: none"> – Treatment/strategy A is recommended/indicated in preference to treatment B – Treatment A should be chosen over treatment B 	LEVEL A <ul style="list-style-type: none"> • High-quality evidence‡ from more than 1 RCT • Meta-analyses of high-quality RCTs • One or more RCTs corroborated by high-quality registry studies
CLASS 2a (MODERATE) Benefit >> Risk Suggested phrases for writing recommendations: <ul style="list-style-type: none"> • Is reasonable • Can be useful/effective/beneficial • Comparative-Effectiveness Phrases‡: <ul style="list-style-type: none"> – Treatment/strategy A is probably recommended/indicated in preference to treatment B – It is reasonable to choose treatment A over treatment B 	LEVEL B-R (Randomized) <ul style="list-style-type: none"> • Moderate-quality evidence‡ from 1 or more RCTs • Meta-analyses of moderate-quality RCTs
CLASS 2b (WEAK) Benefit ≥ Risk Suggested phrases for writing recommendations: <ul style="list-style-type: none"> • May/might be reasonable • May/might be considered • Usefulness/effectiveness is unknown/unclear/uncertain or not well-established 	LEVEL B-NR (Nonrandomized) <ul style="list-style-type: none"> • Moderate-quality evidence‡ from 1 or more well-designed, well-executed nonrandomized studies, observational studies, or registry studies • Meta-analyses of such studies
CLASS 3: No Benefit (MODERATE) Benefit = Risk (Generally, LOE A or B use only) Suggested phrases for writing recommendations: <ul style="list-style-type: none"> • Is not recommended • Is not indicated/useful/effective/beneficial • Should not be performed/administered/other 	LEVEL C-LD (Limited Data) <ul style="list-style-type: none"> • Randomized or nonrandomized observational or registry studies with limitations of design or execution • Meta-analyses of such studies • Physiological or mechanistic studies in human subjects
Class 3: Harm (STRONG) Risk > Benefit Suggested phrases for writing recommendations: <ul style="list-style-type: none"> • Potentially harmful • Causes harm • Associated with excess morbidity/mortality • Should not be performed/administered/other 	LEVEL C-EO (Expert Opinion) <ul style="list-style-type: none"> • Consensus of expert opinion based on clinical experience

COR and LOE are determined independently (any COR may be paired with any LOE).

A recommendation with LOE C does not imply that the recommendation is weak. Many important clinical questions addressed in guidelines do not lend themselves to clinical trials. Although RCTs are unavailable, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

* The outcome or result of the intervention should be specified (an improved clinical outcome or increased diagnostic accuracy or incremental prognostic information).

† For comparative-effectiveness recommendations (COR 1 and 2a; LOE A and B only), studies that support the use of comparator verbs should involve direct comparisons of the treatments or strategies being evaluated.

‡ The method of assessing quality is evolving, including the application of standardized, widely-used, and preferably validated evidence grading tools; and for systematic reviews, the incorporation of an Evidence Review Committee.

COR indicates Class of Recommendation; EO, expert opinion; LD, limited data; LOE, Level of Evidence; NR, nonrandomized; R, randomized; and RCT, randomized controlled trial.

accounting for >365,000 deaths annually (4). Distinguishing between serious and benign causes of chest pain is imperative. The lifetime prevalence of chest pain in the United States is 20% to 40% (5), and women experience this symptom more often than men (6). Of all ED patients with chest pain, only 5.1% will have an acute coronary syndrome (ACS) and more than half will ultimately be found to have a noncardiac cause (7). Nonetheless, chest pain is the most common symptom of CAD in both men and women.

1.1. Document Review and Approval

This document was reviewed by 16 official reviewers nominated by the ACC, AHA, the American College of Emergency Physicians, American Society of Echocardiography (ASE), American Society of Nuclear Cardiology (ASNC), American College of Chest Physicians (CHEST),

Society for Academic Emergency Medicine (SAEM), Society of Cardiovascular Computed Tomography (SCCT), and Society for Cardiovascular Magnetic Resonance (SCMR), and 39 individual content reviewers. Authors' relationships with industry and other entities information is published in [Appendix 1](#) of the full guideline (1). Reviewers' relationships with industry and other entities information is published in [Appendix 2](#) of the full guideline (1).

1.2. Class of Recommendations and Level of Evidence

The Class of Recommendation (COR) indicates the strength of recommendation, encompassing the estimated magnitude and certainty of benefit in proportion to risk. The Level of Evidence (LOE) rates the quality of scientific evidence supporting the intervention on the basis of the type, quantity, and consistency of data from clinical trials and other sources ([Table 1](#)) (8).


1.3. Defining Chest Pain

Figure 2 presents an index of suspicion that chest “pain” is ischemic in origin based on commonly used descriptors.

Recommendations for Defining Chest Pain

Referenced studies that support the recommendations are summarized in [Online Data Supplements 1 and 2](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. An initial assessment of chest pain is recommended to triage patients effectively on the basis of the likelihood that symptoms may be attributable to myocardial ischemia (9-15).
1	C-LD	2. Chest pain should not be described as atypical, because it is not helpful in determining the cause and can be misinterpreted as benign in nature. Instead, chest pain should be described as cardiac, possibly cardiac, or noncardiac because these terms are more specific to the potential underlying diagnosis.




2. INITIAL EVALUATION

2.1. History

Figure 3 presents the top 10 causes of chest pain in ED based on age. Table 2 presents chest pain characteristics and corresponding causes.

Recommendation for History

COR	LOE	RECOMMENDATION
1	C-LD	1. In patients with chest pain, a focused history that includes characteristics and duration of symptoms relative to presentation as well as associated features, and cardiovascular risk factor assessment should be obtained.

FIGURE 3 Top 10 Causes of Chest Pain in the ED Based on Age (Weighted Percentage)

Created using data from Hsia, RY, et al. (7). ED indicates emergency department.

TABLE 2 Chest Pain Characteristics and Corresponding Causes**Nature**

Anginal symptoms are perceived as retrosternal chest discomfort (e.g., pain, discomfort, heaviness, tightness, pressure, constriction, squeezing) (See Section 1.4.2, Defining Chest Pain, in the full guideline [1]).

Sharp chest pain that increases with inspiration and lying supine is unlikely related to ischemic heart disease (e.g., these symptoms usually occur with acute pericarditis).

Onset and duration

Anginal symptoms gradually build in intensity over a few minutes.

Sudden onset of ripping chest pain (with radiation to the upper or lower back) is unlikely to be anginal and is suspicious of an acute aortic syndrome.

Fleeting chest pain—of few seconds' duration—is unlikely to be related to ischemic heart disease.

Location and radiation

Pain that can be localized to a very limited area and pain radiating to below the umbilicus or hip are unlikely related to myocardial ischemia.

Severity

Ripping chest pain ("worse chest pain of my life"), especially when sudden in onset and occurring in a hypertensive patient, or with a known bicuspid aortic valve or aortic dilation, is suspicious of an acute aortic syndrome (e.g., aortic dissection).

Precipitating factors

Physical exercise or emotional stress are common triggers of anginal symptoms.

Occurrence at rest or with minimal exertion associated with anginal symptoms usually indicates ACS.

Positional chest pain is usually nonischemic (e.g., musculoskeletal).

Relieving factors

Relief with nitroglycerin is not necessarily diagnostic of myocardial ischemia and should not be used as a diagnostic criterion.

Associated symptoms

Common symptoms associated with myocardial ischemia include, but are not limited to, dyspnea, palpitations, diaphoresis, lightheadedness, presyncope or syncope, upper abdominal pain, or heartburn unrelated to meals and nausea or vomiting.

Symptoms on the left or right side of the chest, stabbing, sharp pain, or discomfort in the throat or abdomen may occur in patients with diabetes, women, and elderly patients.

ACS indicates acute coronary syndrome.

2.1.1. A Focus on the Uniqueness of Chest Pain in Women

Recommendations for a Focus on the Uniqueness of Chest Pain in Women
 Referenced studies that support the recommendations are summarized in [Online Data Supplements 3 and 4](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. Women who present with chest pain are at risk for underdiagnosis, and potential cardiac causes should always be considered (11,12,14,16-19).
1	B-NR	2. In women presenting with chest pain, it is recommended to obtain a history that emphasizes accompanying symptoms that are more common in women with ACS (11,12,14,16-19).

2.1.2. Considerations for Older Patients With Chest Pain

Recommendation for Considerations for Older Patients With Chest Pain

COR	LOE	RECOMMENDATION
1	C-LD	1. In patients with chest pain who are >75 years of age, ACS should be considered when accompanying symptoms such as shortness of breath, syncope, or acute delirium are present, or when an unexplained fall has occurred (20).

2.1.3. Considerations for Diverse Patient Populations With Chest Pain

Recommendations for Considerations for Diverse Patient Populations With Chest Pain

COR	LOE	RECOMMENDATIONS
1	C-LD	1. Cultural competency training is recommended to help achieve the best outcomes in patients of diverse racial and ethnic backgrounds who present with chest pain.
1	C-LD	2. Among patients of diverse race and ethnicity presenting with chest pain in whom English may not be their primary language, addressing language barriers with the use of formal translation services is recommended.

2.1.4. Patient-Centric Considerations

Recommendation for Patient-Centric Considerations

COR	LOE	RECOMMENDATION
1	C-LD	1. In patients with acute chest pain, it is recommended that 9-1-1 be activated by patients or bystanders to initiate transport to the closest ED by emergency medical services (EMS) (21).

2.2. Physical Examination

Table 3 presents physical examination in patients with chest pain.

Recommendation for Physical Examination

COR	LOE	RECOMMENDATION
1	C-EO	1. In patients presenting with chest pain, a focused cardiovascular examination should be performed initially to aid in the diagnosis of ACS or other potentially serious causes of chest pain (e.g., aortic dissection, pulmonary embolism (PE), or esophageal rupture) and to identify complications.

TABLE 3 Physical Examination in Patients With Chest Pain

Clinical Syndrome	Findings
Emergency	
ACS	Diaphoresis, tachypnea, tachycardia, hypotension, crackles, S3, MR murmur (22); examination may be normal in uncomplicated cases
PE	Tachycardia + dyspnea—>90% of patients; pain with inspiration (23)
Aortic dissection	Connective tissue disorders (e.g., Marfan syndrome), extremity pulse differential (30% of patients, type A>B) (24) Severe pain, abrupt onset + pulse differential + widened mediastinum on CXR >80% probability of dissection (25) Frequency of syncope >10% (24), AR 40%-75% (type A) (26)
Esophageal rupture	Emesis, subcutaneous emphysema, pneumothorax (20% patients), unilateral decreased or absent breath sounds
Other	
Noncoronary cardiac: AS, AR, HCM	AS: Characteristic systolic murmur, tardus or parvus carotid pulse AR: Diastolic murmur at right of sternum, rapid carotid upstroke HCM: Increased or displaced left ventricular impulse, prominent a wave in jugular venous pressure, systolic murmur
Pericarditis	Fever, pleuritic chest pain, increased in supine position, friction rub
Myocarditis	Fever, chest pain, heart failure, S3
Esophagitis, peptic ulcer disease, gall bladder disease	Epigastric tenderness Right upper quadrant tenderness, Murphy sign
Pneumonia	Fever, localized chest pain, may be pleuritic, friction rub may be present, regional dullness to percussion, egophony
Pneumothorax	Dyspnea and pain on inspiration, unilateral absence of breath sounds
Costochondritis, Tietze syndrome	Tenderness of costochondral joints
Herpes zoster	Pain in dermatomal distribution, triggered by touch; characteristic rash (unilateral and dermatomal distribution)

ACS indicates acute coronary syndrome; AR, aortic regurgitation; AS, aortic stenosis; CXR, chest x-ray; LR, likelihood ratio; HCM, hypertrophic cardiomyopathy; MR, mitral regurgitation; PE, pulmonary embolism; and PUD, peptic ulcer disease.

2.3. Diagnostic Testing

2.3.1. Setting Considerations

Recommendations for Setting Considerations

Referenced studies that support the recommendations are summarized in [Online Data Supplement 5](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. Unless a noncardiac cause is evident, an ECG should be performed for patients seen in the office setting with stable chest pain; if an ECG is unavailable the patient should be referred to the ED so one can be obtained (27-31).
1	C-LD	2. Patients with clinical evidence of ACS or other life-threatening causes of acute chest pain seen in the office setting should be transported urgently to the ED, ideally by EMS (27-35).

(continued)


1	C-LD	3. In all patients who present with acute chest pain regardless of the setting, an ECG should be acquired and reviewed for ST-segment-elevation myocardial infarction (STEMI) within 10 minutes of arrival (27-29,32,33,36).
1	C-LD	4. In all patients presenting to the ED with acute chest pain and suspected ACS, cTn should be measured as soon as possible after presentation (34,35).
3: Harm	C-LD	5. For patients with acute chest pain and suspected ACS initially evaluated in the office setting, delayed transfer to the ED for cTn or other diagnostic testing should be avoided.

2.3.2. Electrocardiogram

Figure 4 presents electrocardiographic-directed management of chest pain.

Recommendations for Electrocardiogram (ECG)
 Referenced studies that support the recommendations are summarized in [Online Data Supplement 6](#).

COR	LOE	RECOMMENDATIONS
1	C-EO	1. In patients with chest pain in which an initial ECG is nondiagnostic, serial ECGs to detect potential ischemic changes should be performed, especially when clinical suspicion of ACS is high, symptoms are persistent, or the clinical condition deteriorates (33).
1	C-EO	2. Patients with chest pain in whom the initial ECG is consistent with an ACS should be treated according to STEMI and non-ST-segment-elevation ACS guidelines (32,33).
2a	B-NR	3. In patients with chest pain and intermediate-to-high clinical suspicion for ACS in whom the initial ECG is nondiagnostic, supplemental electrocardiographic leads V ₇ to V ₉ are reasonable to rule out posterior myocardial infarction (MI) (37-39).



2.3.3. Chest Radiography

Recommendation for Chest Radiography

COR	LOE	RECOMMENDATION
1	C-EO	1. In patients presenting with acute chest pain, a chest radiograph is useful to evaluate for other potential cardiac, pulmonary, and thoracic causes of symptoms.

2.3.4. Biomarkers

Recommendations for Biomarkers

Referenced studies that support the recommendations are summarized in [Online Data Supplement 7](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. In patients presenting with acute chest pain, serial cTn I or T levels are useful to identify abnormal values and a rising or falling pattern indicative of acute myocardial injury (35,40-59).
1	B-NR	2. In patients presenting with acute chest pain, high-sensitivity cTn is the preferred biomarker because it enables more rapid detection or exclusion of myocardial injury and increases diagnostic accuracy (35,56,60-63).

(continued)

1	C-EO	3. Clinicians should be familiar with the analytical performance and the 99th percentile upper reference limit that defines myocardial injury for the cTn assay used at their institution (34,61).
3: No benefit	B-NR	4. With availability of cTn, creatine kinase myocardial (CK-MB) isoenzyme and myoglobin are not useful for diagnosis of acute myocardial injury (64-69).

3. CARDIAC TESTING GENERAL CONSIDERATIONS

The approach outlined in this guideline focuses on selective use of testing, optimization of lower cost evaluations,

reducing layered testing, and deferring or eliminating testing when the diagnostic yield is low (Figure 5). Figure 6 display choosing the right diagnostic test. Table 4 presents contraindication by type of imaging modality.





FIGURE 6 Choosing the Right Diagnostic Test



	Favors use of CCTA	Favors use of stress imaging
Goal	<ul style="list-style-type: none"> • Rule out obstructive CAD • Detect nonobstructive CAD 	<ul style="list-style-type: none"> • Ischemia-guided management
Availability and expertise	<ul style="list-style-type: none"> • High-quality imaging and expert interpretation routinely available 	<ul style="list-style-type: none"> • High-quality imaging and expert interpretation routinely available
Likelihood of obstructive CAD	<ul style="list-style-type: none"> • Age <65 y 	<ul style="list-style-type: none"> • Age ≥65 y
Prior test results	<ul style="list-style-type: none"> • Prior functional study inconclusive 	<ul style="list-style-type: none"> • Prior CCTA inconclusive
Other compelling indications	<ul style="list-style-type: none"> • Anomalous coronary arteries • Require evaluation of aorta or pulmonary arteries 	<ul style="list-style-type: none"> • Suspect scar (especially if PET or stress CMR available) • Suspect coronary microvascular dysfunction (when PET or CMR available)

Stress testing information					
	ETT	Stress echocardiography	SPECT MPI	PET MPI	Stress CMR MPI
Patient capable of exercise	✓	✓	✓		
Pharmacologic stress indicated		✓	✓	✓	✓
Quantitative flow				✓	✓
LV dysfunction/scar		✓	✓	✓	✓

ASCVD indicates atherosclerotic cardiovascular disease; CAD, coronary artery disease; CAC, coronary artery calcium; CCTA, coronary computed tomography angiography; CMR, cardiovascular magnetic resonance; ETT, exercise tolerance test; LV, left ventricular; MPI, myocardial perfusion imaging; PET, positron emission tomography; and SPECT, single-photon emission computed tomography.

TABLE 4 Contraindication by Type of Imaging Modality and Stress Protocol

Exercise ECG	Stress Nuclear (70)*	Stress Echocardiography (71-73a)	Stress CMR (74)	CCTA (75)*
<ul style="list-style-type: none"> ■ Abnormal ST changes on resting ECG, digoxin, left bundle branch block, Wolff-Parkinson-White pattern, ventricular paced rhythm (unless test is performed to establish exercise capacity and not for diagnosis of ischemia) ■ Unable to achieve ≥ 5 METs or unsafe to exercise ■ High-risk unstable angina or AMI (<2 d) i.e., active ACS ■ Uncontrolled heart failure ■ Significant cardiac arrhythmias (e.g., VT, complete atrioventricular block) or high risk for arrhythmias caused by QT prolongation ■ Severe symptomatic aortic stenosis ■ Severe systemic arterial hypertension (e.g., $\geq 200/110$ mm Hg) ■ Acute illness (e.g., acute PE, acute myocarditis/pericarditis, acute aortic dissection) 	<ul style="list-style-type: none"> ■ High-risk unstable angina, complicated ACS or AMI (<2 d) ■ Contraindications to vasodilator administration <ul style="list-style-type: none"> ■ Significant arrhythmias (e.g., VT, second- or third-degree atrioventricular block) or sinus bradycardia <45 bpm ■ Significant hypotension (SBP <90 mm Hg) ■ Known or suspected bronchoconstrictive or bronchospastic disease ■ Recent use of dipyridamole or dipyridamole-containing medications ■ Use of methylxanthines (e.g., aminophylline, caffeine) within 12 h ■ Known hypersensitivity to adenosine, regadenoson ■ Severe systemic arterial hypertension (e.g., $\geq 200/110$ mm Hg) 	<ul style="list-style-type: none"> ■ Limited acoustic windows (e.g., in COPD patients) ■ Inability to reach target heart rate ■ Uncontrolled heart failure ■ High-risk unstable angina, active ACS or AMI (<2 d) ■ Serious ventricular arrhythmia or high risk for arrhythmias attributable to QT prolongation ■ Respiratory failure ■ Severe COPD, acute pulmonary emboli, severe pulmonary hypertension ■ Contraindications to dobutamine (if pharmacologic stress test needed) <ul style="list-style-type: none"> ■ Atrioventricular block, uncontrolled atrial fibrillation ■ Critical aortic stenosis† ■ Acute illness (e.g., acute PE, acute myocarditis/pericarditis, acute aortic dissection) ■ Hemodynamically significant LV outflow tract obstruction ■ Contraindications to atropine use: <ul style="list-style-type: none"> ■ Narrow-angle glaucoma ■ Myasthenia gravis ■ Obstructive uropathy ■ Obstructive gastrointestinal disorders ■ Severe systemic arterial hypertension (e.g., $\geq 200/110$ mm Hg) <p><u>Use of Contrast Contraindicated in:</u></p> <ul style="list-style-type: none"> ■ Hypersensitivity to perflutren ■ Hypersensitivity to blood, blood products, or albumin (for Optison only) 	<ul style="list-style-type: none"> ■ Reduced GFR (<30 mL/min/1.73 m²) ■ Contraindications to vasodilator administration ■ Implanted devices not safe for CMR or producing artifact limiting scan quality/interpretation ■ Significant claustrophobia ■ Caffeine use within past 12 h 	<ul style="list-style-type: none"> ■ Allergy to iodinated contrast ■ Inability to cooperate with scan acquisition and/or breath-hold instructions ■ Clinical instability (e.g., acute myocardial infarction, decompensated heart failure, severe hypotension) ■ Renal impairment as defined by local protocols ■ Contraindication to beta blockade in the presence of an elevated heart rate and no alternative medications available for achieving target heart rate ■ Heart rate variability and arrhythmia ■ Contraindication to nitroglycerin (if indicated)

For all the imaging modalities, inability to achieve high-quality images should be considered, in particular for obese patients


*Screening for potential pregnancy by history and/or pregnancy testing should be performed according to the local imaging facilities policies for undertaking radiological examinations that involve ionizing radiation in women of child-bearing age.

†Low-dose dobutamine may be useful for assessing for low-gradient AS.

ACS indicates acute coronary syndrome; AMI, acute myocardial infarction; AS, aortic stenosis; CCTA, coronary computed tomography angiography; CMR, cardiovascular magnetic resonance imaging; COPD, chronic obstructive pulmonary disease; GFR, glomerular filtration rate; LV, left ventricular; MET, metabolic equivalent; MRI, magnetic resonance imaging; PE, pulmonary embolism; SBP, systolic blood pressure; and VT, ventricular tachycardia.

4. CHOOSING THE RIGHT PATHWAY WITH PATIENT-CENTRIC ALGORITHMS FOR ACUTE CHEST PAIN

Figure 7 provides an overview of a patient-centric algorithm for acute chest pain.



4.1. Patients With Acute Chest Pain and Suspected Acute Coronary Syndrome (Not Including STEMI)


Recommendations for Patients With Acute Chest Pain and Suspected ACS (Not Including STEMI)
 Referenced studies that support the recommendations are summarized in [Online Data Supplements 8 and 9](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. In patients presenting with acute chest pain and suspected ACS, clinical decision pathways (CDPs) should categorize patients into low-, intermediate-, and high-risk strata to facilitate disposition and subsequent diagnostic evaluation (40-52,76).
1	B-NR	2. In the evaluation of patients presenting with acute chest pain and suspected ACS for whom serial troponins are indicated to exclude myocardial injury, recommended time intervals after the initial troponin sample collection (time zero) for repeat measurements are: 1 to 3 hours for high-sensitivity troponin and 3 to 6 hours for conventional troponin assays (35,56,77).
1	C-LD	3. To standardize the detection and differentiation of myocardial injury in patients presenting with acute chest pain and suspected ACS, institutions should implement a CDP that includes a protocol for troponin sampling based on their particular assay (78,79).
1	C-LD	4. In patients with acute chest pain and suspected ACS, previous testing when available should be considered and incorporated into CDPs (80-84).
2a	B-NR	5. For patients with acute chest pain, a normal ECG, and symptoms suggestive of ACS that began at least 3 hours before ED arrival, a single hs-cTn concentration that is below the limit of detection on initial measurement (time zero) is reasonable to exclude myocardial injury (51,85-89).

Patients with acute chest pain and suspected ACS cover a spectrum of disease likelihood and stratification into low- versus intermediate- or high-risk groups once STEMI has been excluded (Figure 8). Chest pain risk scores provide a summative assessment combining clinical information, such as age, ST-segment changes on ECG,

symptoms, CAD risk factors, and cTn (Table 5) to estimate a patient’s probability of ACS or risk of 30-day major adverse cardiovascular events (MACE) (90-95). The warranty period of prior cardiac testing should be considered, when symptoms are unchanged (Table 6). Low-risk chest pain has been defined in Table 7.

FIGURE 8 General Approach to Risk Stratification of Patients With Suspected ACS



ACS indicates acute coronary syndrome; CDP, clinical decision pathway; and ECG, electrocardiogram.

TABLE 5 Sample Clinical Decision Pathways Used to Define Risk

	HEART Pathway (91)	EDACS (96)	ADAPT (mADAPT) (97)	NOTR (94)	2020 ESC/hs-cTn* (98,99)	2016 ESC/GRACE (49,100)
Target population	Suspected ACS	Suspected ACS, CP >5 min, planned serial troponin	Suspected ACS, CP >5 min, planned observation	Suspected ACS, ECG, troponin ordered	Suspected ACS, stable	Suspected ACS, planned serial troponin
Target outcome	↑ ED discharge without increasing missed 30-d or 1-y MACE	↑ ED discharge rate without increasing missed 30-d MACE	↑ ED discharge rate without increasing missed 30-d MACE	↑ Low-risk classification without increasing missed 30-d MACE	Early detection of AMI; 30-d MACE	Early detection of AMI
Patients with primary outcome in study population, %	6-22	12	15	5-8	9.8	10-17
Troponin	cTn, hs-cTn	hs-cTn	cTn, hs-cTn	cTn, hs-cTn	hs-cTn	cTn, hs-cTn
Variables used	History ECG Age Risk factors Troponin (0, 3 h)	Age Sex Risk factors History Troponin (0, 2 h)	TIMI score 0-1 No ischemic ECG changes Troponin (0, 2 h)	Age Risk factors Previous AMI or CAD Troponin (0, 2 h)	History ECG hs-cTn (0, 1 or 2 h)	Age HR, SBP Serum Cr Cardiac arrest ECG Cardiac biomarker Killip class
Risk thresholds:						
■ Low risk	HEART score <3 Neg 0, 3-h cTn Neg 0, 2-h hs-cTn	EDACS score <16 Neg 0, 2 h hs-cTn No ischemic ECG Δ	TIMI score 0 (or <1 for mADAPT) ■ Neg 0, 2-h cTn or hs-cTn ■ No ischemic ECG Δ	Age <50 y <3 risk factors Previous AMI or CAD Neg cTn or hs-cTn (0, 2 h)	■ Initial hs-cTn is "very low" and Sx onset >3 h ago Or ■ Initial hs-cTn "low" and 1- or 2-h hs-cTn Δ is "low"	Chest pain free, GRACE <140 ■ Sx <6 h - hs-cTn <ULN (0, 3 h) ■ Sx >6 h - hs-cTn <ULN (arrival)
■ Intermediate risk	HEART score 4-6	NA	TIMI score 2-4	NA	■ Initial hs-cTn is between "low" and "high" And/Or ■ 1- or 2-h hs-cTn Δ is between low and high thresholds	■ TO hs-cTn = 12-52 ng/L or ■ 1-h Δ = 3-5 ng/L
■ High risk	HEART score 7-10 (101,102)	NA	TIMI score 5-7 (102)	NA	■ Initial hs-cTn is "high" Or ■ 1- or 2-h hs-cTn Δ is high	■ TO hs-cTn >52 ng/L or ■ Δ 1 h >5 ng/L
Performance	↑ ED discharges by 21% (40% versus 18%) ↓ 30-d objective testing by 12% (69% versus 57%) ↓ length of stay by 12 h (9.9 versus 21.9 h)	More patients identified as low risk versus ADAPT (42% versus 31%)	ADAPT: More discharged ≤6 h (19% versus 11%)	30-d MACE sensitivity =100% 28% eligible for ED discharge	AMI sensitivity >99% 62% Ruled out (0.2% 30-d MACE) 25% Observe 13% Rule in	AMI sensitivity >99% 30-d MACE not studied
AMI sensitivity, %	100	100	100	100	>99	96.7
cTn accuracy: 30-d MACE sensitivity, %	100	100	100	100	NA	NA
hs-cTn accuracy: 30-d MACE sensitivity, %	95	92	93	99	99	-
ED discharge, %	40	49	19 (ADAPT) 39 (mADAPT)	28	-	-

*The terms "very low," "low," "high," "1 h Δ," and "2 h Δ" refer to hs-cTn assay-specific thresholds published in the ESC guideline (98,99).

ACS indicates acute coronary syndrome; ADAPT, Accelerated Diagnostic protocol to Assess chest Pain using Troponins; AMI, acute myocardial infarction; CP, chest pain or equivalent; Cr, creatinine; cTn, cardiac troponin; hs-cTn, high-sensitivity cardiac troponin; ECG, electrocardiogram; ED, emergency department; EDACS, emergency department ACS; ESC, European Society of Cardiology; GRACE, Global Registry of Acute Coronary Events; HEART, history, ECG, age, risk factors, troponin; HR, heart rate; hs, high sensitivity; MACE, major adverse cardiovascular events; mADAPT, modified (including TIMI scores of 1) ADAPT; NA, not applicable; neg, negative; NICE, National Institute for Health and Clinical Excellence; NOTR, No Objective Testing Rule; SBP, systolic blood pressure; SSACS, symptoms suggestive of ACS; Sx, symptoms; and ULN, upper limit of normal.

TABLE 6 Warranty Period for Prior Cardiac Testing

Test Modality	Result	Warranty Period
Anatomic	Normal coronary angiogram CCTA with no stenosis or plaque	2 y
Stress testing	Normal stress test (given adequate stress)	1 y

CCTA indicates coronary computed tomographic angiography.

TABLE 7 Definition Used for Low-Risk Patients With Chest Pain

Low Risk (<1% 30-d Risk for Death or MACE)	
hs-cTn Based	
T-0	T-0 hs-cTn below the assay limit of detection or "very low" threshold if symptoms present for at least 3 h
T-0 and 1- or 2-h Delta	T-0 hs-cTn and 1- or 2-h delta are both below the assay "low" thresholds (>99% NPV for 30-d MACE)
Clinical Decision Pathway Based	
HEART Pathway (91)	HEART score ≤ 3 , initial and serial cTn/hs-cTn < assay 99th percentile
EDACS (105)	EDACS score ≤ 16 ; initial and serial cTn/hs-cTn < assay 99th percentile
ADAPT (90)	TIMI score 0, initial and serial cTn/hs-cTn < assay 99th percentile
mADAPT	TIMI score 0/1, initial and serial cTn/hs-cTn < assay 99th percentile
NOTR (94)	0 factors

ADAPT indicates 2-hour Accelerated Diagnostic Protocol to Access Patients with Chest Pain Symptoms Using Contemporary Troponins as the Only Biomarkers; cTn, cardiac troponin; EDACS, Emergency Department Acute Coronary Syndrome; HEART Pathway, History, ECG, Age, Risk Factors, Troponin; hs-cTn, high-sensitivity cardiac troponin; MACE, major adverse cardiovascular events; mADAPT, modified 2-hour Accelerated Diagnostic Protocol to Access Patients with Chest Pain Symptoms Using Contemporary Troponins as the Only Biomarkers; NOTR, No Objective Testing Rule; NPV, negative predictive value; and TIMI, Thrombolysis in Myocardial Infarction.

4.1.1. Low-Risk Patients With Acute Chest Pain

Recommendations for Low-Risk Patients With Acute Chest Pain

Referenced studies that support the recommendations are summarized in [Online Data Supplements 10 and 11](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. Patients with acute chest pain and a 30-day risk of death or major adverse cardiovascular events (MACE) <1% should be designated as low risk (34,41,45,49,51,52,54,55,57,85,103).
2a	B-R	2. In patients with acute chest pain and suspected ACS who are deemed low-risk (<1% 30-day risk of death or MACE), it is reasonable to discharge home without admission or urgent cardiac testing (60,94,97,104,105).

4.1.2. Intermediate-Risk Patients With Acute Chest Pain

Recommendations for Intermediate-Risk Patients With Acute Chest Pain
 Referenced studies that support the recommendations are summarized in [Online Data Supplements 12 and 13](#).

COR	LOE	RECOMMENDATIONS
1	C-EO	1. For intermediate-risk patients with acute chest pain, transthoracic echocardiography (TTE) is recommended as a rapid, bedside test to establish baseline ventricular and valvular function, evaluate for wall motion abnormalities, and to assess for pericardial effusion.
2a	A	2. For intermediate-risk patients with acute chest pain, management in an observation unit is reasonable to shorten length of stay and lower cost relative to an inpatient admission (106-112).


4.1.2.1. Intermediate-Risk Patients With Acute Chest Pain and No Known Coronary Artery Disease

Figure 9 presents an evaluation algorithm for patients with suspected ACS at intermediate risk with no known CAD.

Recommendations for Intermediate-Risk Patients With No Known CAD
 Referenced studies that support the recommendations are summarized in [Online Data Supplements 14 and 15](#).

COR	LOE	RECOMMENDATIONS
Index Diagnostic Testing		
Anatomic Testing		
1	A	1. For intermediate-risk patients with acute chest pain and no known CAD eligible for diagnostic testing after a negative or inconclusive evaluation for ACS, coronary computed tomography angiography (CCTA) is useful for exclusion of atherosclerotic plaque and obstructive CAD (113-123).
1	C-EO	2. For intermediate-risk patients with acute chest pain, moderate-severe ischemia on current or prior (≤ 1 year) stress testing, and no known CAD established by prior anatomic testing, invasive coronary angiography (ICA) is recommended.
2a	C-LD	3. For intermediate-risk patients with acute chest pain with evidence of previous mildly abnormal stress test results (≤ 1 year), CCTA is reasonable for diagnosing obstructive CAD (124,125).
Stress Testing		
1	B-NR	4. For intermediate-risk patients with acute chest pain and no known CAD who are eligible for cardiac testing, either exercise ECG, stress echocardiography, stress positron emission tomography (PET)/single-photon emission computed tomography (SPECT) myocardial perfusion imaging (MPI), or stress CMR is useful for the diagnosis of myocardial ischemia (33,107,111,113,116,122,126-145).
Sequential or Add-on Diagnostic Testing		
2a	B-NR	5. For intermediate-risk patients with acute chest pain and no known CAD, with a coronary artery stenosis of 40% to 90% in a proximal or middle coronary artery on CCTA, fractional flow reserve computed tomography (FFR-CT) can be useful for the diagnosis of vessel-specific ischemia and to guide decision-making regarding the use of coronary revascularization (146-152).
2a	C-EO	6. For intermediate-risk patients with acute chest pain and no known CAD, as well as an inconclusive prior stress test, CCTA can be useful for excluding the presence of atherosclerotic plaque and obstructive CAD.
2a	C-EO	7. For intermediate-risk patients with acute chest pain and no known CAD, with an inconclusive CCTA, stress imaging (with echocardiography, PET/SPECT MPI, or CMR) can be useful for the diagnosis of myocardial ischemia.

FIGURE 9 Evaluation Algorithm for Patients With Suspected ACS at Intermediate Risk With No Known CAD



Test choice should be guided by local availability and expertise. *Recent negative test: normal CCTA ≤ 2 years (no plaque/no stenosis) OR negative stress test ≤ 1 year, given adequate stress. †High-risk CAD means left main stenosis $\geq 50\%$; anatomically significant 3-vessel disease ($\geq 70\%$ stenosis). ‡For FFR-CT, turnaround times may impact prompt clinical care decisions. However, the use of FFR-CT does not require additional testing, as would be the case when adding stress testing. CAD indicates coronary artery disease; CCTA, coronary CT angiography; CMR, cardiovascular magnetic resonance imaging; CT, computed tomography; FFR-CT, fractional flow reserve with CT; GDMT, guideline-directed medical therapy; ICA, invasive coronary angiography; INOCA, ischemia and no obstructive coronary artery disease; PET, positron emission tomography; and SPECT, single-photon emission CT.

4.1.2.2. Intermediate-Risk Patients With Acute Chest Pain and Known Coronary Artery Disease


Recommendations for Intermediate-Risk Patients With Acute Chest Pain and Known CAD
Referenced studies that support the recommendations are summarized in [Online Data Supplements 16 and 17](#).

COR	LOE	RECOMMENDATIONS
1	A	1. For intermediate-risk patients with acute chest pain who have known CAD and present with new onset or worsening symptoms, guideline-directed medical therapy (GDMT) should be optimized before additional cardiac testing is performed (153,154).
1	A	2. For intermediate-risk patients with acute chest pain who have worsening frequency of symptoms with significant left main, proximal left anterior descending stenosis, or multivessel CAD on prior anatomic testing or history of prior coronary revascularization, ICA is recommended (113-115,119,155,156).
2a	B-NR	3. For intermediate-risk patients with acute chest pain and known nonobstructive CAD, CCTA can be useful to determine progression of atherosclerotic plaque and obstructive CAD (157-159).

(continued)

- | | | |
|----|------|--|
| 2a | B-NR | 4. For intermediate-risk patients with acute chest pain and coronary artery stenosis of 40% to 90% in a proximal or middle segment on CCTA, FFR-CT is reasonable for diagnosis of vessel-specific ischemia and to guide decision-making regarding the use of coronary revascularization (146,148,149,151,152,160). |
| 2a | B-NR | 5. For intermediate-risk patients with acute chest pain and known CAD who have new onset or worsening symptoms, stress imaging (PET/SPECT MPI, CMR, or stress echocardiography) is reasonable (120,126,129,142). |

Figure 10 includes the evaluation algorithm for patients with known CAD, including patients with nonobstructive and obstructive CAD.



4.1.3. High-Risk Patients With Acute Chest Pain

Recommendations for High-Risk Patients With Acute Chest Pain

Referenced studies that support the recommendations are summarized in [Online Data Supplements 18 and 19](#).

COR	LOE	RECOMMENDATIONS
Recommendations for High-Risk Patients, Including Those With High-Risk Findings on CCTA or Stress Testing		
1	B-NR	1. For patients with acute chest pain and suspected ACS who have new ischemic changes on electrocardiography, troponin-confirmed acute myocardial injury, new-onset left ventricular systolic dysfunction (ejection fraction <40%), newly diagnosed moderate-severe ischemia on stress testing, hemodynamic instability, and/or a high CDP risk score should be designated as high risk for short-term MACE (161-163).
1	C-EO	2. For patients with acute chest pain and suspected ACS who are designated as high risk, ICA is recommended (33,164-166).
2a	B-NR	3. For high-risk patients with acute chest pain who are troponin positive in whom obstructive CAD has been excluded by CCTA or ICA, CMR or echocardiography can be effective in establishing alternative diagnoses (167-171).

4.1.4. Acute Chest Pain in Patients With Prior Coronary Artery Bypass Graft (CABG) Surgery

Recommendations for Acute Chest Pain in Patients With Prior CABG Surgery

COR	LOE	RECOMMENDATIONS
1	C-LD	1. In patients with prior CABG surgery presenting with acute chest pain who do not have ACS, performing stress imaging is effective to evaluate for myocardial ischemia or CCTA for graft stenosis or occlusion (172-178).
1	C-LD	2. In patients with prior CABG surgery presenting with acute chest pain, who do not have ACS (165,179-184) or who have an indeterminate/nondiagnostic stress test, ICA is useful (179).

4.1.5. Evaluation of Patients With Acute Chest Pain Receiving Dialysis

Recommendation for Evaluation of Patients With Acute Chest Pain Receiving Dialysis

Referenced studies that support the recommendation are summarized in [Online Data Supplement 20](#).

COR	LOE	RECOMMENDATION
1	B-NR	1. In patients who experience acute unremitting chest pain while undergoing dialysis, transfer by EMS to an acute care setting is recommended (185-189).

4.1.6. Evaluation of Acute Chest Pain in Patients With Cocaine and Methamphetamine Use

Recommendation for Evaluation of Acute Chest Pain in Patients With Cocaine and Methamphetamine Use

Referenced studies that support the recommendation are summarized in [Online Data Supplement 21](#).

COR	LOE	RECOMMENDATION
2a	B-NR	1. In patients presenting with acute chest pain, it is reasonable to consider cocaine and methamphetamine use as a cause of their symptoms (190-192).

4.1.7. Shared Decision-Making in Patients With Acute Chest Pain

Recommendations for Shared Decision-Making in Patients With Acute Chest Pain
 Referenced studies that support the recommendations are summarized in [Online Data Supplement 22](#).

COR	LOE	RECOMMENDATIONS
1	B-R	1. For patients with acute chest pain and suspected ACS who are deemed low risk by a CDP, patient decision aids are beneficial to improve understanding and effectively facilitate risk communication (193,194).
1	B-R	2. For patients with acute chest pain and suspected ACS who are deemed intermediate risk by a CDP, shared decision-making between the clinician and patient regarding the need for admission, for observation, discharge, or further evaluation in an outpatient setting is recommended for improving patient understanding and reducing low-value testing (193,194).

4.2. Evaluation of Acute Chest Pain With Nonischemic Cardiac Pathologies

Recommendation for Evaluation of Acute Chest Pain With Nonischemic Cardiac Pathologies

COR	LOE	RECOMMENDATION
1	C-EO	1. In patients with acute chest pain in whom other potentially life-threatening nonischemic cardiac conditions are suspected (e.g., aortic pathology, pericardial effusion, endocarditis), TTE is recommended for diagnosis.

4.2.1. Acute Chest Pain With Suspected Acute Aortic Syndrome

Recommendations for Acute Chest Pain With Suspected Acute Aortic Syndrome

COR	LOE	RECOMMENDATIONS
1	C-EO	1. In patients with acute chest pain where there is clinical concern for aortic dissection, computed tomography angiography (CTA) of the chest, abdomen, and pelvis is recommended for diagnosis and treatment planning.
1	C-EO	2. In patients with acute chest pain where there is clinical concern for aortic dissection, transesophageal echocardiography (TEE) or CMR should be performed to make the diagnosis if CT is contraindicated or unavailable.

4.2.2. Acute Chest Pain With Suspected PE

Recommendations for Acute Chest Pain With Suspected PE
 Referenced studies that support the recommendations are summarized in [Online Data Supplement 23](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. In stable patients with acute chest pain with high clinical suspicion for PE, CTA using a PE protocol is recommended (195-198).
1	C-EO	2. For patients with acute chest pain and possible PE, need for further testing should be guided by pretest probability.

4.2.3. Acute Chest Pain With Suspected Myopericarditis

Recommendations for Acute Chest Pain With Suspected Myopericarditis
Referenced studies that support the recommendations are summarized in [Online Data Supplement 24](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. In patients with acute chest pain and myocardial injury who have nonobstructive coronary arteries on anatomic testing, CMR with gadolinium contrast is effective to distinguish myopericarditis from other causes, including myocardial infarction and nonobstructive coronary arteries (MINOCA) (168,170,171,199-201).
1	B-NR	2. In patients with acute chest pain with suspected acute myopericarditis, CMR is useful if there is diagnostic uncertainty, or to determine the presence and extent of myocardial and pericardial inflammation and fibrosis (202-207).
1	C-EO	3. In patients with acute chest pain and suspected myopericarditis, TTE is effective to determine the presence of ventricular wall motion abnormalities, pericardial effusion, valvular abnormalities, or restrictive physiology.
2b	C-LD	4. In patients with acute chest pain with suspected acute pericarditis, noncontrast or contrast cardiac CT scanning may be reasonable to determine the presence and degree of pericardial thickening (202,203,208).

4.2.4. Acute Chest Pain With Valvular Heart Disease (VHD)

Recommendations for Acute Chest Pain With VHD

COR	LOE	RECOMMENDATIONS
1	C-EO	1. In patients presenting with acute chest pain with suspected or known history of VHD, TTE is useful in determining the presence, severity, and cause of VHD.
1	C-EO	2. In patients presenting with acute chest pain with suspected or known VHD in whom TTE diagnostic quality is inadequate, TEE (with 3D imaging if available) is useful in determining the severity and cause of VHD.
2a	C-EO	3. In patients presenting with acute chest pain with known or suspected VHD, CMR imaging is reasonable as an alternative to TTE and/or TEE is nondiagnostic.

4.3. Evaluation of Acute Chest Pain With Suspected Noncardiac Causes

Recommendation for Evaluation of Acute Chest Pain With Suspected Noncardiac Causes

COR	LOE	RECOMMENDATION
1	C-EO	1. Patients with acute chest pain should be evaluated for noncardiac causes if they have persistent or recurring symptoms despite a negative stress test or anatomic cardiac evaluation, or a low-risk designation by a CDP.

The differential diagnosis for noncardiac causes of acute chest pain is quite broad, and includes respiratory, musculoskeletal, gastrointestinal, psychological, and other causes (Table 8).

TABLE 8 Differential Diagnosis of Noncardiac Chest Pain

Respiratory
Pulmonary embolism
Pneumothorax/hemothorax
Pneumomediastinum
Pneumonia
Bronchitis
Pleural irritation
Malignancy
Gastrointestinal
Cholecystitis
Pancreatitis
Hiatal hernia
Gastroesophageal reflux disease/gastritis/esophagitis
Peptic ulcer disease
Esophageal spasm
Dyspepsia
Chest wall
Costochondritis
Chest wall trauma or inflammation
Herpes zoster (shingles)
Cervical radiculopathy
Breast disease
Rib fracture
Musculoskeletal injury/spasm
Psychological
Panic disorder
Anxiety
Clinical depression
Somatization disorder
Hypochondria
Other
Hyperventilation syndrome
Carbon monoxide poisoning
Sarcoidosis
Lead poisoning
Prolapsed intervertebral disc
Thoracic outlet syndrome
Adverse effect of certain medications (e.g., 5-fluorouracil)
Sickle cell crisis

4.3.1. Evaluation of Acute Chest Pain With Suspected Gastrointestinal Syndromes

Recommendation for Evaluation of Acute Chest Pain With Suspected Gastrointestinal Syndromes

COR	LOE	RECOMMENDATION
2a	C-LD	1. In patients with recurrent acute chest pain without evidence of a cardiac or pulmonary cause, evaluation for gastrointestinal causes is reasonable.

4.3.2. Evaluation of Acute Chest Pain With Suspected Anxiety and Other Psychosomatic Considerations

Recommendation for Evaluation of Acute Chest Pain With Suspected Anxiety and Other Psychosomatic Considerations Referenced studies that support the recommendation are summarized in [Online Data Supplement 25](#).

COR	LOE	RECOMMENDATION
2a	B-R	1. For patients with recurrent, similar presentations for acute chest pain with no evidence of a physiological cause on prior diagnostic evaluation including a negative workup for myocardial ischemia, referral to a cognitive-behavioral therapist is reasonable (209-222).

4.3.3. Evaluation of Acute Chest Pain in Patients With Sickle Cell Disease

Recommendations for Evaluation of Acute Chest Pain in Patients With Sickle Cell Disease Referenced studies that support the recommendations are summarized in [Online Data Supplement 26](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. In patients with sickle cell disease who report acute chest pain, emergency transfer by EMS to an acute care setting is recommended (223-227).
1	C-LD	2. In patients with sickle cell disease who report acute chest pain, ACS should be excluded (225-227).

5. EVALUATION OF PATIENTS WITH STABLE CHEST PAIN


5.1. Patients With No Known CAD Presenting With Stable Chest Pain

Stable chest pain is a symptom of myocardial ischemia characterized by chest pain that is provoked with stress (physical or emotional). Risk status in stable ischemic heart disease (SIHD) is not well defined. [Figure 11](#) provides a description of SIHD risk estimates (228).

FIGURE 11 Pretest Probabilities of Obstructive CAD in Symptomatic Patients According to Age, Sex, and Symptoms

Pretest Probabilities of Obstructive CAD in Symptomatic Patients
 (A) according to age, sex, and symptoms;
 (B) according to age, sex, symptoms, and CAC

Age, y	Chest Pain		Dyspnea	
	Men	Women	Men	Women
30-39	≤4	≤5	0	3
40-49	≤22	≤10	12	3
50-59	≤32	≤13	20	9
60-69	≤44	≤16	27	14
70+	≤52	≤27	32	12



Modified from Juarez-Orozco, et al. (228) and Winther S, et al. (229). 1) The pretest probability shown is for patients with anginal symptoms. Patients with lower-risk symptoms would be expected to have lower pretest probability. 2) The darker green- and orange-shaded regions denote the groups in which noninvasive testing is most beneficial (pretest probability >15%). The light green-shaded regions denote the groups with pretest probability of CAD ≤15% in which the testing for diagnosis may be considered based on clinical judgment (228). 3) If CAC is available, it can also be used to estimate the pretest probability based on CAC score (229). CAC indicates coronary artery calcium; and CAD, coronary artery disease.

5.1.2. Low-Risk Patients With Stable Chest Pain and No Known CAD

Recommendations for Low-Risk Patients With Stable Chest Pain and No Known CAD
 Referenced studies that support the recommendations are summarized in [Online Data Supplements 27 and 28](#).

COR	LOE	RECOMMENDATIONS
1	B-NR	1. For patients with stable chest pain and no known CAD presenting to the outpatient clinic, a model to estimate pretest probability of obstructive CAD is effective to identify patients at low risk for obstructive CAD and favorable prognosis in whom additional diagnostic testing can be deferred (228-232).
2a	B-R	2. For patients with stable chest pain and no known CAD categorized as low risk, CAC testing is reasonable as a first-line test for excluding calcified plaque and identifying patients with a low likelihood of obstructive CAD (233-236).
2a	B-NR	3. For patients with stable chest pain and no known CAD categorized as low risk, exercise testing without imaging is reasonable as a first-line test for excluding myocardial ischemia and determining functional capacity in patients with an interpretable ECG (237).


5.1.3. Intermediate-High Risk Patients With Stable Chest Pain and No Known CAD

Figure 12 presents a CDP for patients with stable chest pain and no known CAD.

Recommendations for Intermediate-High Risk Patients With Stable Chest Pain and No Known CAD Referenced studies that support the recommendations are summarized in Online Data Supplements 29 and 30.

COR	LOE	RECOMMENDATIONS
Index Diagnostic Testing		
Anatomic Testing		
1	A	1. For intermediate-high risk patients with stable chest pain and no known CAD, CCTA is effective for diagnosis of CAD, for risk stratification, and for guiding treatment decisions (160,238-248).
Stress Testing		
1	B-R	2. For intermediate-high risk patients with stable chest pain and no known CAD, stress imaging (stress echocardiography, PET/SPECT MPI or CMR) is effective for diagnosis of myocardial ischemia and for estimating risk of MACE (124,245,249-270).
2a	B-R	3. For intermediate-high risk patients with stable chest pain and no known CAD for whom rest/stress nuclear MPI is selected, PET is reasonable in preference to SPECT, if available to improve diagnostic accuracy and decrease the rate of nondiagnostic test results (271-274).
2a	B-R	4. For intermediate-high risk patients with stable chest pain and no known CAD with an interpretable ECG and ability to achieve maximal levels of exercise (≥ 5 metabolic equivalent [MET]s), exercise electrocardiography is reasonable (181,237,245,249,251,275-278).
2b	B-NR	5. In intermediate-high risk patients with stable chest pain selected for stress MPI using SPECT, the use of attenuation correction or prone imaging may be reasonable to decrease the rate of false-positive findings (279-284).
Assessment of Left Ventricular Function		
1	B-NR	6. In intermediate-high risk patients with stable chest pain who have pathological Q waves, symptoms or signs suggestive of heart failure, complex ventricular arrhythmias, or a heart murmur with unclear diagnosis, use of TTE is effective for diagnosis of resting left ventricular systolic and diastolic ventricular function and detection of myocardial, valvular, and pericardial abnormalities (249,250,285).
Sequential or Add-on Testing: What to Do If Index Test Results Are Positive or Inconclusive		
2a	B-NR	7. For intermediate-high risk patients with stable chest pain and known coronary stenosis of 40% to 90% in a proximal or middle coronary segment on CCTA, FFR-CT can be useful for diagnosis of vessel-specific ischemia and to guide decision-making regarding the use of coronary revascularization (146,148,149,160,286-288).
2a	B-NR	8. For intermediate-high risk patients with stable chest pain after an inconclusive or abnormal exercise ECG or stress imaging study, CCTA is reasonable (84,154,242,289-291).
2a	B-NR	9. For intermediate-high risk patients with stable chest pain and no known CAD undergoing stress testing, the addition of CAC testing can be useful (235,292-297).
2a	B-NR	10. For intermediate-high risk patients with stable chest pain after inconclusive CCTA, stress imaging is reasonable (237,249,250,255-258,298-303).
2b	C-EO	11. For intermediate-high risk patients with stable chest pain after a negative stress test but with high clinical suspicion of CAD, CCTA or ICA may be reasonable.

FIGURE 12 Clinical Decision Pathway for Patients With Stable Chest Pain and No Known CAD



Test choice should be guided by local availability and expertise. *Test choice guided by patient's exercise capacity, resting electrocardiographic abnormalities; CCTA preferable in those <65 years of age and not on optimal preventive therapies; stress testing favored in those ≥65 years of age (with a higher likelihood of ischemia). †High-risk CAD means left main stenosis ≥50%; anatomically significant 3-vessel disease (≥70% stenosis). CAD indicates coronary artery disease; CCTA, coronary CT angiography; CMR, cardiovascular magnetic resonance imaging; CT, computed tomography; FFR-CT, fractional flow reserve with CT; GDMT, guideline-directed medical therapy; INOCA, ischemia and no obstructive CAD; PET, positron emission tomography; and SPECT, single-photon emission CT.

5.2. Patients With Known CAD Presenting With Stable Chest Pain

Recommendations for Patients With Known CAD Presenting With Stable Chest Pain

Referenced studies that support the recommendations are summarized in [Online Data Supplement 31](#).

COR	LOE	RECOMMENDATIONS
1	A	1. For patients with obstructive CAD and stable chest pain, it is recommended to optimize GDMT (153,154,304).
1	C-EO	2. For patients with known nonobstructive CAD and stable chest pain, it is recommended to optimize preventive therapies (305,306).

5.2.1. Patients With Obstructive CAD Who Present With Stable Chest Pain


Recommendations for Patients With Obstructive CAD Who Present With Stable Chest Pain

Referenced studies that support the recommendations are summarized in [Online Data Supplements 32 and 33](#).

COR	LOE	RECOMMENDATIONS
Index Diagnostic Testing		
Anatomic Testing		
1	A	1. For patients with obstructive CAD who have stable chest pain despite GDMT and moderate-severe ischemia, ICA is recommended for guiding therapeutic decision-making (153,154,304,307).
1	A	2. For patients with obstructive CAD who have stable chest pain despite optimal GDMT, those referred for ICA without prior stress testing benefit from FFR or instantaneous wave free ratio (307-310).
1	B-R	3. For symptomatic patients with obstructive CAD who have stable chest pain with CCTA-defined $\geq 50\%$ stenosis in the left main coronary artery, obstructive CAD with FFR with CT ≤ 0.80 , or severe stenosis ($\geq 70\%$) in all 3 main vessels, ICA is effective for guiding therapeutic decision-making (154,165).
2a	B-NR	4. For patients who have stable chest pain with previous coronary revascularization, CCTA is reasonable to evaluate bypass graft or stent patency (for stents ≥ 3 mm) (288,311-314).
Stress Testing		
1	B-NR	5. For patients with obstructive CAD who have stable chest pain despite optimal GDMT, stress PET/SPECT MPI, CMR, or echocardiography is recommended for diagnosis of myocardial ischemia, estimating risk of MACE, and guiding therapeutic decision-making (265,272,315-335).
2a	B-R	6. For patients with obstructive CAD who have stable chest pain despite optimal GDMT, when selected for rest/stress nuclear MPI, PET is reasonable in preference to SPECT, if available, to improve diagnostic accuracy and decrease the rate of nondiagnostic test results (261).
2a	B-NR	7. For patients with obstructive CAD who have stable chest pain despite GDMT, exercise treadmill testing can be useful to determine if the symptoms are consistent with angina pectoris, assess the severity of symptoms, evaluate functional capacity and select management, including cardiac rehabilitation (154,336-338).
2a	B-NR	8. For patients with obstructive CAD who have stable chest pain symptoms undergoing stress PET MPI or stress CMR, the addition of myocardial blood flow reserve is useful to improve diagnosis accuracy and enhance risk stratification (272,331-335).

Imaging should be considered in those with new onset or persistent stable chest pain (Figure 13).

FIGURE 13 Clinical Decision Pathway for Patients With Stable Chest Pain (or Equivalent) Symptoms With Prior MI, Prior Revascularization, or Known CAD on Invasive Coronary Angiography or CCTA, Including Those With Nonobstructive CAD



Test choice should be guided by local availability and expertise. *Known CAD means prior MI, revascularization, known obstructive CAD, nonobstructive CAD. †High-risk CAD means left main stenosis ≥50%; or obstructive CAD with FFR-CT ≤0.80. ‡Test choice guided by the patient's exercise capacity, resting electrocardiographic abnormalities. §Patients with prior CABG or stents >3.0 mm. *Follow-up Testing and Intensification of GDMT Guided by Initial Test Results and Persistence / Worsening / Frequency of Symptoms and Shared Decision Making.* CABG indicates coronary artery bypass graft; CAD, coronary artery disease; CCTA, coronary CT angiography; CMR, cardiovascular magnetic resonance imaging; CT, computed tomography; ECG, electrocardiogram; FFR-CT, fractional flow reserve with CT; GDMT, guideline-directed medical therapy; ICA, invasive coronary angiography; IFR, instant wave-free ratio; INOCA, ischemia and no obstructive coronary artery disease; MI, myocardial infarction; MPI, myocardial perfusion imaging; PET, positron emission tomography; SIHD, stable ischemic heart disease; and SPECT, single-photon emission CT.

5.2.1.1. Patients With Prior CABG Surgery With Stable Chest Pain

Recommendations for Patients With Prior CABG Surgery With Stable Chest Pain

COR	LOE	RECOMMENDATIONS
1	C-LD	1. In patients who have had prior CABG surgery presenting with stable chest pain whose noninvasive stress test results show moderate-to-severe ischemia (165,179-184), or in those suspected to have myocardial ischemia with indeterminate/nondiagnostic stress test, ICA is recommended for guiding therapeutic decision-making (179).
2a	C-LD	2. In patients who have had prior CABG surgery presenting with stable chest pain who are suspected to have myocardial ischemia, it is reasonable to perform stress imaging or CCTA to evaluate for myocardial ischemia or graft stenosis or occlusion (172-178,339).

5.2.2. Patients With Known Nonobstructive CAD Presenting With Stable Chest Pain

Recommendations for Patients With Known Nonobstructive CAD Presenting With Stable Chest Pain Referenced studies that support the recommendations are summarized in [Online Data Supplements 34 and 35](#).


COR	LOE	RECOMMENDATIONS
Index Diagnostic Testing		
Anatomic Testing		
2a	B-NR	1. For symptomatic patients with known nonobstructive CAD who have stable chest pain, CCTA is reasonable for determining atherosclerotic plaque burden and progression to obstructive CAD, and guiding therapeutic decision-making (124,158,159,340-343).
2a	B-NR	2. For patients with known coronary stenosis from 40% to 90% on CCTA, FFR can be useful for diagnosis of vessel-specific ischemia and to guide decision-making regarding the use of ICA (146,148,149,160,286-288).
Stress Testing		
2a	C-LD	3. For patients with known extensive nonobstructive CAD with stable chest pain symptoms, stress imaging (PET/SPECT, CMR, or echocardiography) is reasonable for the diagnosis of myocardial ischemia (272,328,331-334,344-347).

5.2.3. Patients With Suspected Ischemia and No Obstructive CAD (INOCA)

Recommendations for Patients With Suspected INOCA
 Referenced studies that support the recommendations are summarized in [Online Data Supplements 36 and 37](#).

COR	LOE	RECOMMENDATIONS
2a	B-NR	1. For patients with persistent stable chest pain and nonobstructive CAD and at least mild myocardial ischemia on imaging, it is reasonable to consider invasive coronary function testing to improve the diagnosis of coronary microvascular dysfunction and to enhance risk stratification (348-351).
2a	B-NR	2. For patients with persistent stable chest pain and nonobstructive CAD, stress PET MPI with myocardial blood flow reserve is reasonable to diagnose microvascular dysfunction and enhance risk stratification (272,331-334,344,345).
2a	B-NR	3. For patients with persistent stable chest pain and nonobstructive CAD, stress CMR with the addition of myocardial blood flow reserve measurement is reasonable to improve diagnosis of coronary myocardial dysfunction and for estimating risk of MACE (328,346,347).
2b	C-EO	4. For patients with persistent stable chest pain and nonobstructive CAD, stress echocardiography with the addition of coronary flow velocity reserve measurement may be reasonable to improve diagnosis of coronary myocardial dysfunction and for estimating risk of MACE.

A proposed diagnostic evaluation pathway is outlined in **Figure 14**.



REFERENCES

- Gulati M, Levy PD, Mukherjee D, et al. 2021 AHA/ACC/ASE/CHEST/SAEM/SCCT/SCMR guideline for the evaluation and diagnosis of chest pain: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2021;78:e187-e285.
- Lawton JS, Tamis-Holland JE, Bangalore S, et al. 2021 ACC/AHA/SCAI guideline for coronary artery revascularization: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2021. In press.
- Rui P, Kang K. National Hospital Ambulatory Medical Care Survey: 2017 emergency department summary tables. National Center for Health Statistics. Available at: https://www.cdc.gov/nchs/data/nhamcs/web_tables/2017_ed_web_tables-508.pdf. Accessed February 12, 2021.

3. Rui P, Okeyode T. National Ambulatory Medical Care Survey: 2016 national summary tables. 2016. Available at: https://www.cdc.gov/nchs/data/ahcd/namcs_summary/2016_namcs_web_tables.pdf. Accessed February 12, 2021.
4. Virani SS, Alonso A, Benjamin EJ, et al. Heart disease and stroke statistics-2020 update: a report from the American Heart Association. *Circulation*. 2020;141:e139-e596.
5. Ruigomez A, Rodriguez LA, Wallander MA, et al. Chest pain in general practice: incidence, comorbidity and mortality. *Fam Pract*. 2006;23:167-174.
6. Bosner S, Becker A, Haasensitter J, et al. Chest pain in primary care: epidemiology and pre-work-up probabilities. *Eur J Gen Pract*. 2009;15:141-146.
7. Hsia RY, Hale Z, Tabas JA. A national study of the prevalence of life-threatening diagnoses in patients with chest pain. *JAMA Intern Med*. 2016;176:1029-1032.
8. ACCF/AHA Task Force on Practice Guidelines. Methodology manual and policies from the ACCF/AHA Task Force on Practice Guidelines. American College of Cardiology and American Heart Association. 2010. Available at: <https://www.acc.org/Guidelines/About-Guidelines-and-Clinical-Documents/Methodology> and https://professional.heart.org/-/media/phd-files/guidelines-and-statements/methodology_manual_and_policies_ucm_319826.pdf. Accessed July 26, 2021.
9. Canto JG, Rogers WJ, Goldberg RJ, et al. Association of age and sex with myocardial infarction symptom presentation and in-hospital mortality. *JAMA*. 2012;307:813-822.
10. van der Meer MG, Backus BE, van der Graaf Y, et al. The diagnostic value of clinical symptoms in women and men presenting with chest pain at the emergency department, a prospective cohort study. *PLoS One*. 2015;10:e0116431.
11. Hernal K, Pagidipati NJ, Coles A, et al. Sex differences in demographics, risk factors, presentation, and noninvasive testing in stable outpatients with suspected coronary artery disease: insights from the PROMISE trial. *J Am Coll Cardiol Img*. 2016;9:337-346.
12. Lichtman JH, Leifheit EC, Safdar B, et al. Sex differences in the presentation and perception of symptoms among young patients with myocardial infarction: evidence from the VIRGO Study (Variation in Recovery: Role of Gender on Outcomes of Young AMI Patients). *Circulation*. 2018;137:781-790.
13. Bosner S, Haasensitter J, Hani MA, et al. Gender differences in presentation and diagnosis of chest pain in primary care. *BMC Fam Pract*. 2009;10:79.
14. Ferry AV, Anand A, Strachan FE, et al. Presenting symptoms in men and women diagnosed with myocardial infarction using sex-specific criteria. *J Am Heart Assoc*. 2019;8:e012307.
15. Kreatsoulas C, Fleegler EW, Kubzansky LD, et al. Young adults and adverse childhood events: a potent measure of cardiovascular risk. *Am J Med*. 2019;132:605-613.
16. Garcia M, Mulvagh SL, Merz CN, et al. Cardiovascular disease in women: clinical perspectives. *Circ Res*. 2016;118:1273-1293.
17. Pelletier R, Khan NA, Cox J, et al. Sex versus gender-related characteristics: which predicts outcome after acute coronary syndrome in the young? *J Am Coll Cardiol*. 2016;67:127-135.
18. Kreatsoulas C, Dinakar D, Mehta S, et al. Machine learning to evaluate gender differences in typical and atypical angina among patients with obstructive coronary artery disease. ESC Congress 2019. Paris, France. 2019.
19. DeFilippis EM, Collins BL, Singh A, et al. Women who experience a myocardial infarction at a young age have worse outcomes compared with men: the Mass General Brigham YOUNG-MI registry. *Eur Heart J*. 2020;41:4127-4137.
20. Grosmaître P, Le Vavasour O, Yachouh E, et al. Significance of atypical symptoms for the diagnosis and management of myocardial infarction in elderly patients admitted to emergency departments. *Arch Cardiovasc Dis*. 2013;106:586-592.
21. Leifheit-Limson EC, D'Onofrio G, Daneshvar M, et al. Sex differences in cardiac risk factors, perceived risk, and health care provider discussion of risk and risk modification among young patients with acute myocardial infarction: the VIRGO study. *J Am Coll Cardiol*. 2015;66:1949-1957.
22. McConaghy JR, Oza RS. Outpatient diagnosis of acute chest pain in adults. *Am Fam Physician*. 2013;87:177-182.
23. Perrier A, Desmarais S, Miron MJ, et al. Non-invasive diagnosis of venous thromboembolism in outpatients. *Lancet*. 1999;353:190-195.
24. Tsai TT, Trimarchi S, Nienaber CA. Acute aortic dissection: perspectives from the International Registry of Acute Aortic Dissection (IRAD). *Eur J Vasc Endovasc Surg*. 2009;37:149-159.
25. von Kodolitsch Y, Schwartz AG, Nienaber CA. Clinical prediction of acute aortic dissection. *Arch Intern Med*. 2000;160:2977-2982.
26. Klompas M. Does this patient have an acute thoracic aortic dissection? *JAMA*. 2002;287:2262-2272.
27. Diercks DB, Kontos MC, Chen AY, et al. Utilization and impact of pre-hospital electrocardiograms for patients with acute ST-segment elevation myocardial infarction: data from the NCDR (National Cardiovascular Data Registry) ACTION (Acute Coronary Treatment and Intervention Outcomes Network) Registry. *J Am Coll Cardiol*. 2009;53:161-166.
28. Puymirat E, Simon T, Steg PG, et al. Association of changes in clinical characteristics and management with improvement in survival among patients with ST-elevation myocardial infarction. *JAMA*. 2012;308:998-1006.
29. Jollis JG, Granger CB, Henry TD, et al. Systems of care for ST-segment-elevation myocardial infarction: a report from the American Heart Association's Mission: Lifeline. *Circ Cardiovasc Qual Outcomes*. 2012;5:423-428.
30. Postma S, Bergmeijer T, ten Berg J, et al. Pre-hospital diagnosis, triage and treatment in patients with ST elevation myocardial infarction. *Heart*. 2012;98:1674-1678.
31. Patel M, Dunford JV, Aguilar S, et al. Pre-hospital electrocardiography by emergency medical personnel: effects on scene and transport times for chest pain and ST-segment elevation myocardial infarction patients. *J Am Coll Cardiol*. 2012;60:806-811.
32. O'Gara PT, Kushner FG, Ascheim DD, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;61:e78-e140.
33. Amsterdam EA, Wenger NK, Brindis RG, et al. 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2014;64:e139-e228.
34. Neumann JT, Twerenbold R, Ojeda F, et al. Application of high-sensitivity troponin in suspected myocardial infarction. *N Engl J Med*. 2019;380:2529-2540.
35. Thygesen K, Alpert JS, Jaffe AS, et al. Fourth universal definition of myocardial infarction (2018). *J Am Coll Cardiol*. 2018;72:2231-2264.
36. Becker L, Larsen MP, Eisenberg MS. Incidence of cardiac arrest during self-transport for chest pain. *Ann Emerg Med*. 1996;28:612-616.
37. Vogiatzis I, Koulouris E, Ioannidis A, et al. The importance of the 15-lead versus 12-lead ECG recordings in the diagnosis and treatment of right ventricle and left ventricle posterior and lateral wall acute myocardial infarctions. *Acta Inform Med*. 2019;27:35-39.
38. Ashida T, Tani S, Nagao K, et al. Usefulness of synthesized 18-lead electrocardiography in the diagnosis of ST-elevation myocardial infarction: a pilot study. *Am J Emerg Med*. 2017;35:448-457.
39. Matetzky S, Freimark D, Feinberg MS, et al. Acute myocardial infarction with isolated ST-segment elevation in posterior chest leads V7-9: "hidden" ST-segment elevations revealing acute posterior infarction. *J Am Coll Cardiol*. 1999;34:748-753.
40. Apple FS, Jesse RL, Newby LK, et al. National Academy of Clinical Biochemistry and IFCC Committee for Standardization of Markers of Cardiac Damage Laboratory Medicine Practice Guidelines: analytical issues for biochemical markers of acute coronary syndromes. *Circulation*. 2007;115:e352-e355.
41. Bandstein N, Ljung R, Johansson M, et al. Undetectable high-sensitivity cardiac troponin T level in the emergency department and risk of myocardial infarction. *J Am Coll Cardiol*. 2014;63:2569-2578.
42. Body R, Burrows G, Carley S, et al. High-sensitivity cardiac troponin concentrations below the limit of detection to exclude acute myocardial infarction: a prospective evaluation. *Clin Chem*. 2015;61:983-989.
43. Body R, Carley S, McDowell G, et al. Rapid exclusion of acute myocardial infarction in patients with undetectable troponin using a high-sensitivity assay. *J Am Coll Cardiol*. 2011;58:1332-1339.
44. Boeddinghaus J, Nestelberger T, Twerenbold R, et al. Direct comparison of 4 very early rule-out strategies for acute myocardial infarction using high-sensitivity cardiac troponin I. *Circulation*. 2017;135:1597-1611.
45. Chapman AR, Anand A, Boeddinghaus J, et al. Comparison of the efficacy and safety of early rule-out pathways for acute myocardial infarction. *Circulation*. 2017;135:1586-1596.
46. de Lemos JA, Morrow DA, deFilippi CR. Highly sensitive troponin assays and the cardiology community: a love/hate relationship? *Clin Chem*. 2011;57:826-829.
47. Jaffe AS, Apple FS, Morrow DA, et al. Being rational about (im)precision: a statement from the

- Biochemistry Subcommittee of the Joint European Society of Cardiology/American College of Cardiology Foundation/American Heart Association/World Heart Federation Task Force for the Definition of Myocardial Infarction. *Clin Chem*. 2010;56:941–943.
48. Morrow DA, Cannon CP, Jesse RL, et al. National Academy of Clinical Biochemistry laboratory medicine practice guidelines: clinical characteristics and utilization of biochemical markers in acute coronary syndromes. *Circulation*. 2007;115:e356–e375.
49. Mueller C, Giannitsis E, Christ M, et al. Multicenter evaluation of a 0-hour/1-hour algorithm in the diagnosis of myocardial infarction with high-sensitivity cardiac troponin T. *Ann Emerg Med*. 2016;68:76–87, e4.
50. Odqvist M, Andersson PO, Tygesen H, et al. High-sensitivity troponins and outcomes after myocardial infarction. *J Am Coll Cardiol*. 2018;71:2616–2624.
51. Peacock WF, Baumann BM, Bruton D, et al. Efficacy of high-sensitivity troponin T in identifying very-low-risk patients with possible acute coronary syndrome. *JAMA Cardiol*. 2018;3:104–111.
52. Reichlin T, Schindler C, Drexler B, et al. One-hour rule-out and rule-in of acute myocardial infarction using high-sensitivity cardiac troponin T. *Arch Intern Med*. 2012;172:1211–1218.
53. Reichlin T, Twerenbold R, Maushart C, et al. Risk stratification in patients with unstable angina using absolute serial changes of 3 high-sensitive troponin assays. *Am Heart J*. 2013;165:371–378.e3.
54. Rubini Gimenez M, Hoeller R, Reichlin T, et al. Rapid rule out of acute myocardial infarction using undetectable levels of high-sensitivity cardiac troponin. *Int J Cardiol*. 2013;168:3896–3901.
55. Rubini Gimenez M, Twerenbold R, Jaeger C, et al. One-hour rule-in and rule-out of acute myocardial infarction using high-sensitivity cardiac troponin I. *Am J Med*. 2015;128:861–870, e4.
56. Twerenbold R, Boeddinghaus J, Nestelberger T, et al. Clinical use of high-sensitivity cardiac troponin in patients with suspected myocardial infarction. *J Am Coll Cardiol*. 2017;70:996–1012.
57. Twerenbold R, Neumann JT, Sorensen NA, et al. Prospective validation of the 0/1-h algorithm for early diagnosis of myocardial infarction. *J Am Coll Cardiol*. 2018;72:620–632.
58. Wildi K, Nelles B, Twerenbold R, et al. Safety and efficacy of the 0 h/3 h protocol for rapid rule out of myocardial infarction. *Am Heart J*. 2016;181:16–25.
59. Zhelev Z, Hyde C, Youngman E, et al. Diagnostic accuracy of single baseline measurement of Elecsys Troponin T high-sensitive assay for diagnosis of acute myocardial infarction in emergency department: systematic review and meta-analysis. *BMJ*. 2015;350:h15.
60. Cullen L, Mueller C, Parsonage WA, et al. Validation of high-sensitivity troponin I in a 2-hour diagnostic strategy to assess 30-day outcomes in emergency department patients with possible acute coronary syndrome. *J Am Coll Cardiol*. 2013;62:1242–1249.
61. Lipinski MJ, Baker NC, Escarcega RO, et al. Comparison of conventional and high-sensitivity troponin in patients with chest pain: a collaborative meta-analysis. *Am Heart J*. 2015;169:6–16, e6.
62. Twerenbold R, Wildi K, Jaeger C, et al. Optimal cutoff levels of more sensitive cardiac troponin assays for the early diagnosis of myocardial infarction in patients with renal dysfunction. *Circulation*. 2015;131:2041–2050.
63. van Wijk S, Jacobs L, Eurlings LW, et al. Troponin T measurements by high-sensitivity vs conventional assays for risk stratification in acute dyspnea. *Clin Chem*. 2012;58:284–292.
64. Aviles RJ, Wright RS, Aviles JM, et al. Long-term prognosis of patients with clinical unstable angina pectoris without elevation of creatine kinase but with elevation of cardiac troponin I levels. *Am J Cardiol*. 2002;90:875–878.
65. Eggers KM, Oldgren J, Nordenskjold A, et al. Diagnostic value of serial measurement of cardiac markers in patients with chest pain: limited value of adding myoglobin to troponin I for exclusion of myocardial infarction. *Am Heart J*. 2004;148:574–581.
66. Kavsak PA, MacRae AR, Newman AM, et al. Effects of contemporary troponin assay sensitivity on the utility of the early markers myoglobin and CKMB isoforms in evaluating patients with possible acute myocardial infarction. *Clin Chim Acta*. 2007;380:213–216.
67. Kontos MC, de Lemos JA, Ou FS, et al. Troponin-positive, MB-negative patients with non-ST-elevation myocardial infarction: an undertreated but high-risk patient group: results from the National Cardiovascular Data Registry Acute Coronary Treatment and Intervention Outcomes Network-Get With The Guidelines (NCDR ACTION-GWTG) Registry. *Am Heart J*. 2010;160:819–825.
68. Newby LK, Roe MT, Chen AY, et al. Frequency and clinical implications of discordant creatine kinase-MB and troponin measurements in acute coronary syndromes. *J Am Coll Cardiol*. 2006;47:312–318.
69. Volz KA, McGillicuddy DC, Horowitz GL, et al. Creatine kinase-MB does not add additional benefit to a negative troponin in the evaluation of chest pain. *Am J Emerg Med*. 2012;30:188–190.
70. Henzlova MJ, Duvall WL, Einstein AJ, et al. ASNC imaging guidelines for SPECT nuclear cardiology procedures: stress, protocols, and tracers. *J Nucl Cardiol*. 2016;23:606–639.
71. Senior R, Becher H, Monaghan M, et al. Contrast echocardiography: evidence-based recommendations by European Association of Echocardiography. *Eur J Echocardiogr*. 2009;10:194–212.
72. Patil HR, Main M. Revisiting the safety profile of echocardiography contrast agents. Available at: <https://www.acc.org/latest-in-cardiology/articles/2016/06/23/08/23/revisiting-the-safety-profile-of-echocardiography-contrast-agents>. Accessed August 16, 2020.
73. Porter TR, Abdelmoneim S, Belcik JT, et al. Guidelines for the cardiac sonographer in the performance of contrast echocardiography: a focused update from the American Society of Echocardiography. *J Am Soc Echocardiogr*. 2014;27:797–810.
- 73 a. Pellikka PA, Arruda-Olson A, Chaudhry FA, et al. Guidelines for performance, interpretation, and application of stress echocardiography in ischemic heart disease: from the American Society of Echocardiography. *J Am Soc Echocardiogr*. 2020;33:1–41.e8.
74. Kramer CM, Barkhausen J, Bucciarelli-Ducci C, et al. Standardized cardiovascular magnetic resonance imaging (CMR) protocols: 2020 update. *J Cardiovasc Magn Reson*. 2020;22:17.
75. Abbara S, Blanke P, Maroules CD, et al. SCCT guidelines for the performance and acquisition of coronary computed tomographic angiography: a report of the Society of Cardiovascular Computed Tomography Guidelines Committee. *J Cardiovasc Comput Tomogr*. 2016;10:435–449.
76. Jaffe AS. Chasing troponin: how low can you go if you can see the rise? *J Am Coll Cardiol*. 2006;48:1763–1764.
77. Morrow DA. Clinician's guide to early rule-out strategies with high-sensitivity cardiac troponin. *Circulation*. 2017;135:1612–1616.
78. Januzzi JL Jr, Mahler SA, Christenson RH, et al. Recommendations for institutions transitioning to high-sensitivity troponin testing: JACC Scientific Expert Panel. *J Am Coll Cardiol*. 2019;73:1059–1077.
79. Huis In 't Veld MA, Cullen L, Mahler SA, et al. The fast and the furious: low-risk chest pain and the rapid rule-out protocol. *West J Emerg Med*. 2017;18:474–478.
80. Schulman-Marcus J, Hartaigh BO, Gransar H, et al. Sex-specific associations between coronary artery plaque extent and risk of major adverse cardiovascular events: the CONFIRM Long-Term Registry. *J Am Coll Cardiol Img*. 2016;9:364–372.
81. Hadamitzky M, Freissmuth B, Meyer T, et al. Prognostic value of coronary computed tomographic angiography for prediction of cardiac events in patients with suspected coronary artery disease. *J Am Coll Cardiol Img*. 2009;2:404–411.
82. Finck T, Hardenberg J, Will A, et al. Ten-year follow-up after coronary computed tomography angiography in patients with suspected coronary artery disease. *J Am Coll Cardiol Img*. 2019;12:1330–1338.
83. Hochman JS, Reynolds HR, Bangalore S, et al. Baseline characteristics and risk profiles of participants in the ISCHEMIA randomized clinical trial. *JAMA Cardiol*. 2019;4:273–286.
84. ISCHEMIA Trial Research Group, Maron DJ, Hochman JS, et al. International Study of Comparative Health Effectiveness with Medical and Invasive Approaches (ISCHEMIA) trial: rationale and design. *Am Heart J*. 2018;201:124–135.
85. Pickering JW, Than MP, Cullen L, et al. Rapid rule-out of acute myocardial infarction with a single high-sensitivity cardiac troponin T measurement below the limit of detection: a collaborative meta-analysis. *Ann Intern Med*. 2017;166:715–724.
86. McRae AD, Innes G, Graham M, et al. Undetectable concentrations of a Food and Drug Administration-approved high-sensitivity cardiac troponin T assay to rule out acute myocardial infarction at emergency department arrival. *Acad Emerg Med*. 2017;24:1267–1277.
87. Sandoval Y, Nowak R, deFilippi CR, et al. Myocardial infarction risk stratification with a single measurement of high-sensitivity troponin I. *J Am Coll Cardiol*. 2019;74:271–282.
88. Bularga A, Lee KK, Stewart S, et al. High-sensitivity troponin and the application of risk stratification thresholds in patients with suspected acute

- coronary syndrome. *Circulation*. 2019;140:1557-1568.
- 89.** Chew DP, Lambrakis K, Blyth A, et al. A randomized trial of a 1-hour troponin T protocol in suspected acute coronary syndromes: the Rapid Assessment of Possible Acute Coronary Syndrome in the Emergency Department With High-Sensitivity Troponin T Study (RAPID-TnT). *Circulation*. 2019;140:1543-1556.
- 90.** Than M, Cullen L, Aldous S, et al. 2-Hour accelerated diagnostic protocol to assess patients with chest pain symptoms using contemporary troponins as the only biomarker: the ADAPT trial. *J Am Coll Cardiol*. 2012;59:2091-2098.
- 91.** Mahler SA, Riley RF, Hiestand BC, et al. The HEART Pathway randomized trial: identifying emergency department patients with acute chest pain for early discharge. *Circ Cardiovasc Qual Outcomes*. 2015;8:195-203.
- 92.** Mark DG, Huang J, Chettipally U, et al. Performance of coronary risk scores among patients with chest pain in the emergency department. *J Am Coll Cardiol*. 2018;71:606-616.
- 93.** Stopyra JP, Miller CD, Hiestand BC, et al. Chest pain risk stratification: a comparison of the 2-Hour Accelerated Diagnostic Protocol (ADAPT) and the HEART Pathway. *Crit Pathw Cardiol*. 2016;15:46-49.
- 94.** Stopyra JP, Miller CD, Hiestand BC, et al. Validation of the no objective testing rule and comparison to the HEART Pathway. *Acad Emerg Med*. 2017;24:1165-1168.
- 95.** Stopyra JP, Riley RF, Hiestand BC, et al. The HEART Pathway randomized controlled trial one-year outcomes. *Acad Emerg Med*. 2019;26:41-50.
- 96.** Than MP, Pickering JW, Aldous SJ, et al. Effectiveness of EDACS versus ADAPT accelerated diagnostic pathways for chest pain: a pragmatic randomized controlled trial embedded within practice. *Ann Emerg Med*. 2016;68:93-102, e1.
- 97.** Than M, Aldous S, Lord SJ, et al. A 2-hour diagnostic protocol for possible cardiac chest pain in the emergency department: a randomized clinical trial. *JAMA Intern Med*. 2014;174:51-58.
- 98.** Collet JP, Thiele H, Barbato E, et al. 2020 ESC guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J*. 2021;42:1289-1367.
- 99.** Twerenbold R, Costabel JP, Nestelberger T, et al. Outcome of applying the ESC 0/1-hour algorithm in patients with suspected myocardial infarction. *J Am Coll Cardiol*. 2019;74:483-494.
- 100.** Roffi M, Patrono C, Collet JP, et al. 2015 ESC guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting Without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2016;37:267-315.
- 101.** Backus BE, Six AJ, Kelder JC, et al. A prospective validation of the HEART score for chest pain patients at the emergency department. *Int J Cardiol*. 2013;168:2153-2158.
- 102.** Fanaroff AC, Rymer JA, Goldstein SA, et al. Does this patient with chest pain have acute coronary syndrome?: The rational clinical examination systematic review. *JAMA*. 2015;314:1955-1965.
- 103.** Reichlin T, Twerenbold R, Wildi K, et al. Prospective validation of a 1-hour algorithm to rule-out and rule-in acute myocardial infarction using a high-sensitivity cardiac troponin T assay. *CMAJ*. 2015;187:E243-E252.
- 104.** Greenslade JH, Carlton EW, Van Hise C, et al. Diagnostic accuracy of a new high-sensitivity troponin I assay and five accelerated diagnostic pathways for ruling out acute myocardial infarction and acute coronary syndrome. *Ann Emerg Med*. 2018;71:439-451, e3.
- 105.** Flaws D, Than M, Scheuermeyer FX, et al. External validation of the emergency department assessment of chest pain score accelerated diagnostic pathway (EDACS-ADP). *Emerg Med J*. 2016;33:618-625.
- 106.** Farkouh ME, Smars PA, Reeder GS, et al. A clinical trial of a chest-pain observation unit for patients with unstable angina. Chest Pain Evaluation in the Emergency Room (CHEER) Investigators. *N Engl J Med*. 1998;339:1882-1888.
- 107.** Miller CD, Hwang W, Hoekstra JW, et al. Stress cardiac magnetic resonance imaging with observation unit care reduces cost for patients with emergent chest pain: a randomized trial. *Ann Emerg Med*. 2010;201056:209-219.
- 108.** Roberts RR, Zalenski RJ, Mensah EK, et al. Costs of an emergency department-based accelerated diagnostic protocol vs hospitalization in patients with chest pain: a randomized controlled trial. *JAMA*. 1997;278:1670-1676.
- 109.** Ross MA, Hockenberry JM, Mutter R, et al. Protocol-driven emergency department observation units offer savings, shorter stays, and reduced admissions. *Health Aff (Millwood)*. 2013;32:2149-2156.
- 110.** Rydman RJ, Zalenski RJ, Roberts RR, et al. Patient satisfaction with an emergency department chest pain observation unit. *Ann Emerg Med*. 1997;29:109-115.
- 111.** Miller CD, Hwang W, Case D, et al. Stress CMR imaging observation unit in the emergency department reduces 1-year medical care costs in patients with acute chest pain: a randomized study for comparison with inpatient care. *J Am Coll Cardiol*. 2011;4:862-870.
- 112.** Hockenberry JM, Mutter R, Barrett M, et al. Factors associated with prolonged observation services stays and the impact of long stays on patient cost. *Health Serv Res*. 2014;49:893-909.
- 113.** Goldstein JA, Chinnaiyan KM, Abidov A, et al. The CT-STAT (Coronary Computed Tomographic Angiography for Systematic Triage of Acute Chest Pain Patients to Treatment) trial. *J Am Coll Cardiol*. 2011;58:1414-1422.
- 114.** Litt HI, Gatsonis C, Snyder B, et al. CT angiography for safe discharge of patients with possible acute coronary syndromes. *N Engl J Med*. 2012;366:1393-1403.
- 115.** Goodacre S, Thokala P, Carroll C, et al. Systematic review, meta-analysis and economic modelling of diagnostic strategies for suspected acute coronary syndrome. *Health Technol Assess*. 2013;17. v-vi, 1-188.
- 116.** Goldstein JA, Gallagher MJ, O'Neill WW, et al. A randomized controlled trial of multi-slice coronary computed tomography for evaluation of acute chest pain. *J Am Coll Cardiol*. 2007;49:863-871.
- 117.** Chang SA, Choi SI, Choi EK, et al. Usefulness of 64-slice multidetector computed tomography as an initial diagnostic approach in patients with acute chest pain. *Am Heart J*. 2008;156:375-383.
- 118.** Miller AH, Pepe PE, Peshock R, et al. Is coronary computed tomography angiography a resource sparing strategy in the risk stratification and evaluation of acute chest pain? Results of a randomized controlled trial. *Acad Emerg Med*. 2011;18:458-467.
- 119.** Hoffmann U, Truong QA, Schoenfeld DA, et al. Coronary CT angiography versus standard evaluation in acute chest pain. *N Engl J Med*. 2012;367:299-308.
- 120.** Linde JJ, Kofoed KF, Sorgaard M, et al. Cardiac computed tomography guided treatment strategy in patients with recent acute-onset chest pain: results from the randomised, controlled trial: CARDiac cT in the treatment of acute CHEst pain (CATCH). *Int J Cardiol*. 2013;168:5257-5262.
- 121.** Hamilton-Craig C, Fifoot A, Hansen M, et al. Diagnostic performance and cost of CT angiography versus stress ECG—a randomized prospective study of suspected acute coronary syndrome chest pain in the emergency department (CT-COMPARE). *Int J Cardiol*. 2014;177:867-873.
- 122.** Levsky JM, Spevack DM, Travin MI, et al. Coronary computed tomography angiography versus radionuclide myocardial perfusion imaging in patients with chest pain admitted to telemetry: a randomized trial. *Ann Intern Med*. 2015;163:174-183.
- 123.** Dedic A, Lubbers MM, Schaap J, et al. Coronary CT angiography for suspected ACS in the era of high-sensitivity troponins: randomized multicenter study. *J Am Coll Cardiol*. 2016;67:16-26.
- 124.** Hoffmann U, Ferencik M, Udelson JE, et al. Prognostic value of noninvasive cardiovascular testing in patients with stable chest pain: insights from the PROMISE trial (Prospective Multicenter Imaging Study for Evaluation of Chest Pain). *Circulation*. 2017;135:2320-2332.
- 125.** Min JK, Dunning A, Lin FY, et al. Age- and sex-related differences in all-cause mortality risk based on coronary computed tomography angiography findings results from the International Multicenter CONFIRM (Coronary CT Angiography Evaluation for Clinical Outcomes: An International Multicenter Registry) of 23,854 patients without known coronary artery disease. *J Am Coll Cardiol*. 2011;58:849-860.
- 126.** Blankstein R, Ahmed W, Bamberg F, et al. Comparison of exercise treadmill testing with cardiac computed tomography angiography among patients presenting to the emergency room with chest pain: the Rule Out Myocardial Infarction Using Computer-Assisted Tomography (ROMICAT) study. *Circ Cardiovasc Imaging*. 2012;5:233-242.
- 127.** Jeetley P, Burden L, Stoykova B, et al. Clinical and economic impact of stress echocardiography compared with exercise electrocardiography in patients with suspected acute coronary syndrome but negative troponin: a prospective randomized controlled study. *Eur Heart J*. 2007;28:204-211.
- 128.** Dadkhah S, Almuwaqqat Z, Sulaiman S, et al. Sensitive troponin I and stress testing in the emergency department for the early management of chest pain using 2-hour protocol. *Crit Pathw Cardiol*. 2017;16:89-92.

- 129.** Nucifora G, Badano LP, Sarraf-Zadegan N, et al. Comparison of early dobutamine stress echocardiography and exercise electrocardiographic testing for management of patients presenting to the emergency department with chest pain. *Am J Cardiol.* 2007;100:1068–1073.
- 130.** Jasani G, Papas M, Patel AJ, et al. Immediate stress echocardiography for low-risk chest pain patients in the emergency department: a prospective observational cohort study. *J Emerg Med.* 2018;54:156–164.
- 131.** Krishnan S, Venn R, Blumenthal DM, et al. Utilization of stress testing for low-risk patients with chest discomfort in the emergency department. *J Nucl Cardiol.* 2019;26:1642–1646.
- 132.** Hermann LK, Newman DH, Pleasant WA, et al. Yield of routine provocative cardiac testing among patients in an emergency department-based chest pain unit. *JAMA Intern Med.* 2013;173:1128–1133.
- 133.** Diercks DB, Mumma BE, Frank Peacock W, et al. Incremental value of objective cardiac testing in addition to physician impression and serial contemporary troponin measurements in women. *Acad Emerg Med.* 2013;20:265–270.
- 134.** Poldervaart JM, Six AJ, Backus BE, et al. The predictive value of the exercise ECG for major adverse cardiac events in patients who presented with chest pain in the emergency department. *Clin Res Cardiol.* 2013;102:305–312.
- 135.** Napoli AM, Tran S, Wang J. Low-risk chest pain patients younger than 40 years do not benefit from admission and stress testing. *Crit Pathw Cardiol.* 2013;12:201–203.
- 136.** Scott AC, Bilesky J, Lamanna A, et al. Limited utility of exercise stress testing in the evaluation of suspected acute coronary syndrome in patients aged less than 40 years with intermediate risk features. *Emerg Med Australas.* 2014;26:170–176.
- 137.** Aldous S, Richards AM, Cullen L, et al. The incremental value of stress testing in patients with acute chest pain beyond serial cardiac troponin testing. *Emerg Med J.* 2016;33:319–324.
- 138.** Greenslade JH, Parsonage W, Ho A, et al. Utility of routine exercise stress testing among intermediate risk chest pain patients attending an emergency department. *Heart Lung Circ.* 2015;24:879–884.
- 139.** Skoien W. Diagnostic yield of routine stress testing in low and intermediate risk chest pain patients under 40 years: a systematic review. *Crit Pathw Cardiol.* 2016;15:114–120.
- 140.** Levsky JM, Haramati LB, Spevack DM, et al. Coronary computed tomography angiography versus stress echocardiography in acute chest pain: a randomized controlled trial. *J Am Coll Cardiol Img.* 2018;11:1288–1297.
- 141.** Heitner JF, Klem I, Rasheed D, et al. Stress cardiac MR imaging compared with stress echocardiography in the early evaluation of patients who present to the emergency department with intermediate-risk chest pain. *Radiology.* 2014;271:56–64.
- 142.** Udelson JE, Beshansky JR, Ballin DS, et al. Myocardial perfusion imaging for evaluation and triage of patients with suspected acute cardiac ischemia: a randomized controlled trial. *JAMA.* 2002;288:2693–2700.
- 143.** Miller CD, Case LD, Little WC, et al. Stress CMR reduces revascularization, hospital readmission, and recurrent cardiac testing in intermediate-risk patients with acute chest pain. *J Am Coll Cardiol Img.* 2013;6:785–794.
- 144.** Ingkanisorn WP, Kwong RY, Bohme NS, et al. Prognosis of negative adenosine stress magnetic resonance in patients presenting to an emergency department with chest pain. *J Am Coll Cardiol.* 2006;47:1427–1432.
- 145.** Siontis GC, Mavridis D, Greenwood JP, et al. Outcomes of non-invasive diagnostic modalities for the detection of coronary artery disease: network meta-analysis of diagnostic randomised controlled trials. *BMJ.* 2018;360:k504.
- 146.** Norgaard BL, Leipsic J, Gaur S, et al. Diagnostic performance of noninvasive fractional flow reserve derived from coronary computed tomography angiography in suspected coronary artery disease: the NXT trial (Analysis of Coronary Blood Flow Using CT Angiography: Next Steps). *J Am Coll Cardiol.* 2014;63:1145–1155.
- 147.** Sand NPR, Veien KT, Nielsen SS, et al. Prospective comparison of FFR derived from coronary CT angiography with SPECT perfusion imaging in stable coronary artery disease: the ReASSESS study. *J Am Coll Cardiol Img.* 2018;11:1640–1650.
- 148.** Patel MR, Norgaard BL, Fairbairn TA, et al. 1-Year impact on medical practice and clinical outcomes of FFRCT: the ADVANCE registry. *J Am Coll Cardiol Img.* 2020;13:97–105.
- 149.** Fairbairn TA, Nieman K, Akasaka T, et al. Real-world clinical utility and impact on clinical decision-making of coronary computed tomography angiography-derived fractional flow reserve: lessons from the ADVANCE Registry. *Eur Heart J.* 2018;39:3701–3711.
- 150.** Nakanishi R, Osawa K, Ceponiene I, et al. The diagnostic performance of SPECT-MPI to predict functional significant coronary artery disease by fractional flow reserve derived from CCTA (FFRCT): sub-analysis from ACCURACY and VCTO01 studies. *Int J Cardiovasc Imaging.* 2017;33:2067–2072.
- 151.** Chinnaiyan KM, Safian RD, Gallagher ML, et al. Clinical use of CT-derived fractional flow reserve in the emergency department. *J Am Coll Cardiol Img.* 2020;13:452–461.
- 152.** Ferencik M, Lu MT, Mayrhofer T, et al. Non-invasive fractional flow reserve derived from coronary computed tomography angiography in patients with acute chest pain: subgroup analysis of the ROMICAT II trial. *J Cardiovasc Comput Tomogr.* 2019;13:196–202.
- 153.** Boden WE, O'Rourke RA, Teo KK, et al. Optimal medical therapy with or without PCI for stable coronary disease. *N Engl J Med.* 2007;356:1503–1516.
- 154.** Maron DJ, Hochman JS, Reynolds HR, et al. Initial invasive or conservative strategy for stable coronary disease. *N Engl J Med.* 2020;382:1395–1407.
- 155.** Hulten E, Pickett C, Bittencourt MS, et al. Outcomes after coronary computed tomography angiography in the emergency department: a systematic review and meta-analysis of randomized, controlled trials. *J Am Coll Cardiol.* 2013;61:880–892.
- 156.** Hulten E, Pickett C, Bittencourt MS, et al. Meta-analysis of coronary CT angiography in the emergency department. *Eur Heart J Cardiovasc Imaging.* 2013;14:607.
- 157.** Puchner SB, Liu T, Mayrhofer T, et al. High-risk plaque detected on coronary CT angiography predicts acute coronary syndromes independent of significant stenosis in acute chest pain: results from the ROMICAT-II trial. *J Am Coll Cardiol.* 2014;64:684–692.
- 158.** Motoyama S, Ito H, Sarai M, et al. Plaque characterization by coronary computed tomography angiography and the likelihood of acute coronary events in mid-term follow-up. *J Am Coll Cardiol.* 2015;66:337–346.
- 159.** Chang HJ, Lin FY, Lee SE, et al. Coronary atherosclerotic precursors of acute coronary syndromes. *J Am Coll Cardiol.* 2018;71:2511–2522.
- 160.** Douglas PS, Pontone G, Hlatky MA, et al. Clinical outcomes of fractional flow reserve by computed tomographic angiography-guided diagnostic strategies vs. usual care in patients with suspected coronary artery disease: the prospective longitudinal trial of FFR(CT): outcome and resource impacts study. *Eur Heart J.* 2015;36:3359–3367.
- 161.** Mahmarian JJ, Shaw LJ, Filipchuk NG, et al. A multinational study to establish the value of early adenosine technetium-99m sestamibi myocardial perfusion imaging in identifying a low-risk group for early hospital discharge after acute myocardial infarction. *J Am Coll Cardiol.* 2006;48:2448–2457.
- 162.** Mahmarian JJ, Dakik HA, Filipchuk NG, et al. An initial strategy of intensive medical therapy is comparable to that of coronary revascularization for suppression of scintigraphic ischemia in high-risk but stable survivors of acute myocardial infarction. *J Am Coll Cardiol.* 2006;48:2458–2467.
- 163.** Yan AT, Yan RT, Tan M, et al. Risk scores for risk stratification in acute coronary syndromes: useful but simpler is not necessarily better. *Eur Heart J.* 2007;28:1072–1078.
- 164.** Patel MR, Calhoon JH, Dehmer GJ, et al. 2017 appropriate use criteria for coronary revascularization in patients with stable ischemic heart disease: a report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and Society of Thoracic Surgeons. *J Am Coll Cardiol.* 2017;69:2212–2241.
- 165.** Patel MR, Bailey SR, Bonow RO, et al. ACCF/SCAI/AATS/AHA/ASE/ASNC/HFSA/HRS/SCCM/SCCT/SCMR/STS 2012 appropriate use criteria for diagnostic catheterization: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, Society for Cardiovascular Angiography and Interventions, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol.* 2012;59:1995–2027.

166. Levine GN, Bates ER, Blankenship JC, et al. 2015 ACC/AHA/SCAI focused update on primary percutaneous coronary intervention for patients with ST-elevation myocardial infarction: an update of the 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention and the 2013 ACCF/AHA guideline for the management of ST-Elevation myocardial infarction. *J Am Coll Cardiol*. 2016;67:1235-1250.
167. Pathik B, Raman B, Mohd Amin NH, et al. Troponin-positive chest pain with unobstructed coronary arteries: incremental diagnostic value of cardiovascular magnetic resonance imaging. *Eur Heart J Cardiovasc Imaging*. 2016;17:1146-1152.
168. Dastidar AG, Rodrigues JCL, Johnson TW, et al. Myocardial infarction with nonobstructed coronary arteries: impact of CMR early after presentation. *J Am Coll Cardiol Img*. 2017;10:1204-1206.
169. Assomull RG, Lyne JC, Keenan N, et al. The role of cardiovascular magnetic resonance in patients presenting with chest pain, raised troponin, and unobstructed coronary arteries. *Eur Heart J*. 2007;28:1242-1249.
170. Tornvall P, Gerbaud E, Behaghel A, et al. Myocarditis or "true" infarction by cardiac magnetic resonance in patients with a clinical diagnosis of myocardial infarction without obstructive coronary disease: a meta-analysis of individual patient data. *Atherosclerosis*. 2015;241:87-91.
171. Dastidar AG, Baritussio A, De Garate E, et al. Prognostic role of CMR and conventional risk factors in myocardial infarction with nonobstructed coronary arteries. *J Am Coll Cardiol Img*. 2019;12:1973-1982.
172. Fitzgibbon GM, Kafka HP, Leach AJ, et al. Coronary bypass graft fate and patient outcome: angiographic follow-up of 5,065 grafts related to survival and reoperation in 1,388 patients during 25 years. *J Am Coll Cardiol*. 1996;28:616-626.
173. Harskamp RE, Lopes RD, Baisden CE, et al. Saphenous vein graft failure after coronary artery bypass surgery: pathophysiology, management, and future directions. *Ann Surg*. 2013;257:824-833.
174. Wolk MJ, Bailey SR, Doherty JU, et al. ACCF/AHA/AASE/ASNC/HFSA/HRS/SCAI/SCCT/SCMR/STS 2013 multimodality appropriate use criteria for the detection and risk assessment of stable ischemic heart disease: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. 2014;63:380-406.
175. Taylor AJ, Cerqueira M, Hodgson JM, et al. ACCF/SCCT/ACR/AHA/AASE/ASNC/NASCI/SCAI/SCMR 2010 appropriate use criteria for cardiac computed tomography: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, the Society of Cardiovascular Computed Tomography, the American College of Radiology, the American Heart Association, the American Society of Echocardiography, the American Society of Nuclear Cardiology, the North American Society for Cardiovascular Imaging, the Society for Cardiovascular Angiography and Interventions, and the Society for Cardiovascular Magnetic Resonance. *J Am Coll Cardiol*. 2010;56:1864-1894.
176. Sabik JF 3rd. Understanding saphenous vein graft patency. *Circulation*. 2011;124:273-275.
177. Taggart DP. Current status of arterial grafts for coronary artery bypass grafting. *Ann Cardiothorac Surg*. 2013;2:427-430.
178. Gaudino M, Benedetto U, Fremes S, et al. Radial-artery or saphenous-vein grafts in coronary-artery bypass surgery. *N Engl J Med*. 2018;378:2069-2077.
179. Fihn SD, Blankenship JC, Alexander KP, et al. 2014 ACC/AHA/AATS/PCNA/SCAI/STS focused update of the guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines, and the American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. 2014;64:1929-1949.
180. Hammermeister KE, DeRouen TA, Dodge HT. Variables predictive of survival in patients with coronary disease. Selection by univariate and multivariate analyses from the clinical, electrocardiographic, exercise, arteriographic, and quantitative angiographic evaluations. *Circulation*. 1979;59:421-430.
181. Mark DB, Hlatky MA, Harrell FE Jr, et al. Exercise treadmill score for predicting prognosis in coronary artery disease. *Ann Intern Med*. 1987;106:793-800.
182. Levine GN, Bates ER, Blankenship JC, et al. 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. *J Am Coll Cardiol*. 2011;58:e44-e122.
183. Hillis LD, Smith PK, Anderson JL, et al. 2011 ACCF/AHA guideline for coronary artery bypass graft surgery: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2011;58:e123-e210.
184. Bittl JA, He Y, Jacobs AK, et al. Bayesian methods affirm the use of percutaneous coronary intervention to improve survival in patients with unprotected left main coronary artery disease. *Circulation*. 2013;127:2177-2185.
185. Saran R, Robinson B, Abbott KC, et al. U.S. Renal Data System 2017 Annual Data Report: Epidemiology of Kidney Disease in the United States. *Am J Kidney Dis*. 2018;71:A7.
186. Voroneanu L, Covic A. Arrhythmias in hemodialysis patients. *J Nephrol*. 2009;22:716-725.
187. Herzog CA, Littrell K, Arko C, et al. Clinical characteristics of dialysis patients with acute myocardial infarction in the United States: a collaborative project of the United States Renal Data System and the National Registry of Myocardial Infarction. *Circulation*. 2007;116:1465-1472.
188. Kielstein JT, Abou-Rebyeh F, Hafer C, et al. Right-sided chest pain at the onset of haemodialysis. *Nephrol Dial Transplant*. 2001;16:1493-1495.
189. Modi KS, Gross D, Davidman M. The patient developing chest pain at the onset of haemodialysis sessions—it is not always angina pectoris. *Nephrol Dial Transplant*. 1999;14:221-223.
190. Finkel JB, Marhefka GD. Rethinking cocaine-associated chest pain and acute coronary syndromes. *Mayo Clin Proc*. 2011;86:1198-1207.
191. DeFilippis E, Singh A, Divakaran S, et al. Cocaine and marijuana use among young adults with myocardial infarction. *J Am Coll Cardiol*. 2018;71:2540-2551.
192. Hawley LA, Auten JD, Matteucci MJ, et al. Cardiac complications of adult methamphetamine exposures. *J Emerg Med*. 2013;45:821-827.
193. Hess EP, Hollander JE, Schaffer JT, et al. Shared decision making in patients with low risk chest pain: prospective randomized pragmatic trial. *BMJ*. 2016;355:i6165.
194. Hess EP, Knoedler MA, Shah ND, et al. The chest pain choice decision aid: a randomized trial. *Circ Cardiovasc Qual Outcomes*. 2012;5:251-259.
195. Pollack CV, Schreiber D, Goldhaber SZ, et al. Clinical characteristics, management, and outcomes of patients diagnosed with acute pulmonary embolism in the emergency department: initial report of EMPEROR (Multicenter Emergency Medicine Pulmonary Embolism in the Real World Registry). *J Am Coll Cardiol*. 2011;57:700-706.
196. Konstantinides SV, Torbicki A, Agnelli G, et al. 2014 ESC guidelines on the diagnosis and management of acute pulmonary embolism. *Eur Heart J*. 2014;35:3033-3069, 69a-69k.
197. Raja AS, Greenberg JO, Qaseem A, et al. Evaluation of patients with suspected acute pulmonary embolism: best practice advice from the Clinical Guidelines Committee of the American College of Physicians. *Ann Intern Med*. 2015;163:701-711.
198. Ceriani E, Combesure C, Le Gal G, et al. Clinical prediction rules for pulmonary embolism: a systematic review and meta-analysis. *J Thromb Haemost*. 2010;8:957-970.
199. Bozkurt B, Colvin M, Cook J, et al. Current diagnostic and treatment strategies for specific dilated cardiomyopathies: a scientific statement from the American Heart Association. *Circulation*. 2016;134:e579-e646.
200. Ferreira VM, Schulz-Menger J, Holmvang G, et al. Cardiovascular magnetic resonance in nonischemic myocardial inflammation: expert recommendations. *J Am Coll Cardiol*. 2018;72:3158-3176.
201. Lintings PF, Nivet H, Clement-Guinaudeau S, et al. High-resolution late gadolinium enhancement magnetic resonance for the diagnosis of myocardial infarction with nonobstructed coronary arteries. *J Am Coll Cardiol Img*. 2020;13:1135-1148.
202. Klein AL, Abbara S, Agler DA, et al. American Society of Echocardiography clinical recommendations for multimodality cardiovascular imaging of patients with pericardial disease. *J Am Soc Echocardiogr*. 2013;26:965-1012, e15.
203. Adler Y, Charron P, Imazio M, et al. 2015 ESC guidelines for the diagnosis and management of pericardial diseases: the Task Force for the Diagnosis and Management of Pericardial Diseases of the European Society of Cardiology (ESC). *Eur Heart J*. 2015;36:2921-2964.
204. Taylor AM, Dymarkowski S, Verbeke EK, et al. Detection of pericardial inflammation with late-

- enhancement cardiac magnetic resonance imaging: initial results. *Eur Radiol.* 2006;16:569–574.
- 205.** Young PM, Glockner JF, Williamson EE, et al. MR imaging findings in 76 consecutive surgically proven cases of pericardial disease with CT and pathologic correlation. *Int J Cardiovasc Imaging.* 2012;28:1099–1109.
- 206.** Aquaro GD, Perfetti M, Camastra G, et al. Cardiac MR with late gadolinium enhancement in acute myocarditis with preserved systolic function: ITAMY study. *J Am Coll Cardiol.* 2017;70:1977–1987.
- 207.** Grani C, Buechel RR, Kaufmann PA, et al. Multimodality imaging in individuals with anomalous coronary arteries. *J Am Coll Cardiol Img.* 2017;10:471–481.
- 208.** Hammer MM, Raptis CA, Javidan-Nejad C, et al. Accuracy of computed tomography findings in acute pericarditis. *Acta Radiol.* 2014;55:1197–1202.
- 209.** Carter C, Maddock R, Amsterdam E, et al. Panic disorder and chest pain in the coronary care unit. *Psychosomatics.* 1992;33:302–309.
- 210.** Kline JA, Shapiro NI, Jones AE, et al. Outcomes and radiation exposure of emergency department patients with chest pain and shortness of breath and ultralow pretest probability: a multicenter study. *Ann Emerg Med.* 2014;63:281–288.
- 211.** Webster R, Norman P, Goodacre S, et al. The prevalence and correlates of psychological outcomes in patients with acute non-cardiac chest pain: a systematic review. *Emerg Med J.* 2012;29:267–273.
- 212.** Eslick GD, Talley NJ. Natural history and predictors of outcome for non-cardiac chest pain: a prospective 4-year cohort study. *Neurogastroenterol Motil.* 2008;20:989–997.
- 213.** Foldes-Busque G, Marchand A, Chauny JM, et al. Unexplained chest pain in the ED: could it be panic? *Am J Emerg Med.* 2011;29:743–751.
- 214.** Musey PI Jr, Kline JA. Emergency department cardiopulmonary evaluation of low-risk chest pain patients with self-reported stress and anxiety. *J Emerg Med.* 2017;52:273–279.
- 215.** Al-Ani M, Winchester DE. Prevalence and overlap of noncardiac conditions in the evaluation of low-risk acute chest pain patients. *Crit Pathw Cardiol.* 2015;14:97–102.
- 216.** Czarnecki A, Wang JT, Tu JV, et al. The role of primary care physician and cardiologist follow-up for low-risk patients with chest pain after emergency department assessment. *Am Heart J.* 2014;168:289–295.
- 217.** White KS, Craft JM, Gervino EV. Anxiety and hypervigilance to cardiopulmonary sensations in non-cardiac chest pain patients with and without psychiatric disorders. *Behav Res Ther.* 2010;48:394–401.
- 218.** Burgstaller JM, Jenni BF, Steurer J, et al. Treatment efficacy for non-cardiovascular chest pain: a systematic review and meta-analysis. *PLoS One.* 2014;9:e104722.
- 219.** Kisely SR, Campbell LA, Yelland MJ, et al. Psychological interventions for symptomatic management of non-specific chest pain in patients with normal coronary anatomy. *Cochrane Database Syst Rev.* 2015; CD004101.
- 220.** Mitchell AM, Garvey JL, Chandra A, et al. Prospective multicenter study of quantitative pretest probability assessment to exclude acute coronary syndrome for patients evaluated in emergency department chest pain units. *Ann Emerg Med.* 2006;47:447.
- 221.** Hoorweg BB, Willemsen RT, Cleef LE, et al. Frequency of chest pain in primary care, diagnostic tests performed and final diagnoses. *Heart.* 2017;103:1727–1732.
- 222.** Glombiewski JA, Rief W, Bosner S, et al. The course of nonspecific chest pain in primary care: symptom persistence and health care usage. *Arch Intern Med.* 2010;170:251–255.
- 223.** Platt OS, Brambilla DJ, Rosse WF, et al. Mortality in sickle cell disease. Life expectancy and risk factors for early death. *N Engl J Med.* 1994;330:1639–1644.
- 224.** Thomas AN, Pattison C, Serjeant GR. Causes of death in sickle-cell disease in Jamaica. *Br Med J (Clin Res Ed).* 1982;285:633–635.
- 225.** Vichinsky EP, Styles LA, Colangelo LH, et al. Acute chest syndrome in sickle cell disease: clinical presentation and course. Cooperative Study of Sickle Cell Disease. *Blood.* 1997;89:1787–1792.
- 226.** Martin CR, Johnson CS, Cobb C, et al. Myocardial infarction in sickle cell disease. *J Natl Med Assoc.* 1996;88:428–432.
- 227.** Ogunbayo GO, Misumida N, Olorunfemi O, et al. Comparison of outcomes in patients having acute myocardial infarction with versus without sickle-cell anemia. *Am J Cardiol.* 2017;120:1768–1771.
- 228.** Juarez-Orozco LE, Saraste A, Capodanno D, et al. Impact of a decreasing pre-test probability on the performance of diagnostic tests for coronary artery disease. *Eur Heart J Cardiovasc Imaging.* 2019;20:1198–1207.
- 229.** Winther S, Schmidt SE, Mayrhofer T, et al. Incorporating coronary calcification into pre-test assessment of the likelihood of coronary artery disease. *J Am Coll Cardiol.* 2020;76:2421–2432.
- 230.** Genders TS, Steyerberg EW, Alkadhhi H, et al. A clinical prediction rule for the diagnosis of coronary artery disease: validation, updating, and extension. *Eur Heart J.* 2011;32:1316–1330.
- 231.** Fordyce CB, Douglas PS, Roberts RS, et al. Identification of patients with stable chest pain deriving minimal value from noninvasive testing: the PROMISE minimal-risk tool, a secondary analysis of a randomized clinical trial. *JAMA Cardiol.* 2017;2:400–408.
- 232.** Genders TS, Steyerberg EW, Hunink MG, et al. Prediction model to estimate presence of coronary artery disease: retrospective pooled analysis of existing cohorts. *BMJ.* 2012;344:e3485.
- 233.** Lubbers M, Coenen A, Kofflard M, et al. Comprehensive cardiac CT with myocardial perfusion imaging versus functional testing in suspected coronary artery disease: the multicenter, randomized CRESCENT-II trial. *J Am Coll Cardiol Img.* 2018;11:1625–1636.
- 234.** Lubbers M, Dedic A, Coenen A, et al. Calcium imaging and selective computed tomography angiography in comparison to functional testing for suspected coronary artery disease: the multicenter, randomized CRESCENT trial. *Eur Heart J.* 2016;37:1232–1243.
- 235.** Chang SM, Nabi F, Xu J, et al. The coronary artery calcium score and stress myocardial perfusion imaging provide independent and complementary prediction of cardiac risk. *J Am Coll Cardiol.* 2009;54:1872–1882.
- 236.** Budoff MJ, Mayrhofer T, Ferencik M, et al. Prognostic value of coronary artery calcium in the PROMISE study (Prospective Multicenter Imaging Study for Evaluation of Chest Pain). *Circulation.* 2017;136:1993–2005.
- 237.** Shaw LJ, Mieres JH, Hendel RH, et al. Comparative effectiveness of exercise electrocardiography with or without myocardial perfusion single photon emission computed tomography in women with suspected coronary artery disease: results from the What Is the Optimal Method for Ischemia Evaluation in Women (WOMEN) trial. *Circulation.* 2011;124:1239–1249.
- 238.** Dewey M, Rief M, Martus P, et al. Evaluation of computed tomography in patients with atypical angina or chest pain clinically referred for invasive coronary angiography: randomised controlled trial. *BMJ.* 2016;355:i5441.
- 239.** Meijboom WB, Meijis MF, Schuijff JD, et al. Diagnostic accuracy of 64-slice computed tomography coronary angiography: a prospective, multicenter, multivendor study. *J Am Coll Cardiol.* 2008;52:2135–2144.
- 240.** Budoff MJ, Dowe D, Jollis JG, et al. Diagnostic performance of 64-multidetector row coronary computed tomographic angiography for evaluation of coronary artery stenosis in individuals without known coronary artery disease: results from the prospective multicenter ACCURACY (Assessment by Coronary Computed Tomographic Angiography of Individuals Undergoing Invasive Coronary Angiography) trial. *J Am Coll Cardiol.* 2008;52:1724–1732.
- 241.** Miller JM, Rochitte CE, Dewey M, et al. Diagnostic performance of coronary angiography by 64-row CT. *N Engl J Med.* 2008;359:2324–2336.
- 242.** Chang HJ, Lin FY, Gebow D, et al. Selective referral using CCTA versus direct referral for individuals referred to invasive coronary angiography for suspected CAD: a randomized, controlled, open-label trial. *J Am Coll Cardiol Img.* 2019;12:1303–1312.
- 243.** Sharma A, Coles A, Sekaran NK, et al. Stress testing versus CT angiography in patients with diabetes and suspected coronary artery disease. *J Am Coll Cardiol.* 2019;73:893–902.
- 244.** Min JK, Koduru S, Dunning AM, et al. Coronary CT angiography versus myocardial perfusion imaging for near-term quality of life, cost and radiation exposure: a prospective multicenter randomized pilot trial. *J Cardiovasc Comput Tomogr.* 2012;6:274–283.
- 245.** Douglas PS, Hoffmann U, Patel MR, et al. Outcomes of anatomical versus functional testing for coronary artery disease. *N Engl J Med.* 2015;372:1291–1300.
- 246.** SCOT-HEART Investigators, Newby DE, Adamson PD, et al. Coronary CT angiography and 5-year risk of myocardial infarction. *N Engl J Med.* 2018;379:924–933.
- 247.** SCOT-HEART Investigators. CT coronary angiography in patients with suspected angina due to coronary heart disease (SCOT-HEART): an open-label, parallel-group, multicenter trial. *Lancet.* 2015;385:2383–2391.
- 248.** McKavanagh P, Lusk L, Ball PA, et al. A comparison of cardiac computerized tomography and exercise stress electrocardiogram test for the

investigation of stable chest pain: the clinical results of the CAPP randomized prospective trial. *Eur Heart J Cardiovasc Imaging*. 2015;16:441-448.

249. Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. 2012;60:e44-e164.

250. Douglas PS, Garcia MJ, Haines D, et al. ACCF/AHA/ASNC/HFSA/HRS/SCAI/SCCM/SCCT/SCMR 2011 appropriate use criteria for echocardiography: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Society of Echocardiography, American Heart Association, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, and Society for Cardiovascular Magnetic Resonance. *J Am Coll Cardiol*. 2011;57:1126-1166.

251. Gurunathan S, Zacharias K, Akhtar M, et al. Cost-effectiveness of a management strategy based on exercise echocardiography versus exercise electrocardiography in patients presenting with suspected angina during long term follow up: A randomized study. *Int J Cardiol*. 2018;259:1-7.

252. Carpeggiani C, Landi P, Michelassi C, et al. Stress echocardiography positivity predicts cancer death. *J Am Heart Assoc*. 2017;6:e007104.

253. Abdelmoneim SS, Ball CA, Mantovani F, et al. Prognostic utility of stress testing and cardiac biomarkers in menopausal women at low to intermediate risk for coronary artery disease (SMART study): 5-year outcome. *J Womens Health (Larchmt)*. 2018;27:542-551.

254. Cortigiani L, Urluescu ML, Coltelli M, et al. Apparent declining prognostic value of a negative stress echocardiography based on regional wall motion abnormalities in patients with normal resting left ventricular function due to the changing referral profile of the population under study. *Circ Cardiovasc Imaging*. 2019;12:e008564.

255. Gibbons RJ, Hodge DO, Berman DS, et al. Long-term outcome of patients with intermediate-risk exercise electrocardiograms who do not have myocardial perfusion defects on radionuclide imaging. *Circulation*. 1999;100:2140-2145.

256. Rozanski A, Gansar H, Min JK, et al. Long-term mortality following normal exercise myocardial perfusion SPECT according to coronary disease risk factors. *J Nucl Cardiol*. 2014;21:341-350.

257. Bourque JM, Holland BH, Watson DD, et al. Achieving an exercise workload of > or = 10 metabolic equivalents predicts a very low risk of inducible ischemia: does myocardial perfusion imaging have a role? *J Am Coll Cardiol*. 2009;54:538-545.

258. Shaw LJ, Wilson PW, Hachamovitch R, et al. Improved near-term coronary artery disease risk

classification with gated stress myocardial perfusion SPECT. *J Am Coll Cardiol Img*. 2010;3:1139-1148.

259. Shaw LJ, Min JK, Hachamovitch R, et al. Nomograms for estimating coronary artery disease prognosis with gated stress myocardial perfusion SPECT. *J Nucl Cardiol*. 2012;19:43-52.

260. Uretsky S, Rozanski A. Long-term outcomes following a normal stress myocardial perfusion scan. *J Nucl Cardiol*. 2013;20:715-718.

261. Patel KK, Al Badarin F, Chan PS, et al. Randomized comparison of clinical effectiveness of pharmacologic SPECT and PET MPI in symptomatic CAD patients. *J Am Coll Cardiol Img*. 2019;12:1821-1831.

262. Dorbala S, Di Carli MF, Beanlands RS, et al. Prognostic value of stress myocardial perfusion positron emission tomography: results from a multicenter observational registry. *J Am Coll Cardiol*. 2013;61:176-184.

263. Kay J, Dorbala S, Goyal A, et al. Influence of sex on risk stratification with stress myocardial perfusion Rb-82 positron emission tomography: Results from the PET (Positron Emission Tomography) Prognosis Multicenter Registry. *J Am Coll Cardiol*. 2013;62:1866-1876.

264. Greenwood JP, Maredia N, Younger JF, et al. Cardiovascular magnetic resonance and single-photon emission computed tomography for diagnosis of coronary heart disease (CE-MARC): a prospective trial. *Lancet*. 2012;379:453-460.

265. Schwitler J, Wacker CM, Wilke N, et al. MR-IMPACT II: Magnetic Resonance Imaging for Myocardial Perfusion Assessment in Coronary artery disease Trial: perfusion-cardiac magnetic resonance vs. single-photon emission computed tomography for the detection of coronary artery disease: a comparative multicentre, multivendor trial. *Eur Heart J*. 2013;34:775-781.

266. Hamon M, Fau G, Nee G, et al. Meta-analysis of the diagnostic performance of stress perfusion cardiovascular magnetic resonance for detection of coronary artery disease. *J Cardiovasc Magn Reson*. 2010;12:29.

267. Jaarsma C, Leiner T, Bekkers SC, et al. Diagnostic performance of noninvasive myocardial perfusion imaging using single-photon emission computed tomography, cardiac magnetic resonance, and positron emission tomography imaging for the detection of obstructive coronary artery disease: a meta-analysis. *J Am Coll Cardiol*. 2012;59:1719-1728.

268. Schwitler J, Wacker CM, Wilke N, et al. Superior diagnostic performance of perfusion-cardiovascular magnetic resonance versus SPECT to detect coronary artery disease: the secondary endpoints of the multicenter multivendor MR-IMPACT II (Magnetic Resonance Imaging for Myocardial Perfusion Assessment in Coronary Artery Disease Trial). *J Cardiovasc Magn Reson*. 2012;14:61.

269. Greenwood JP, Ripley DP, Berry C, et al. Effect of care guided by cardiovascular magnetic resonance, myocardial perfusion scintigraphy, or NICE guidelines on subsequent unnecessary angiography rates: the CE-MARC 2 randomized clinical trial. *JAMA*. 2016;316:1051-1060.

270. Nagel E, Greenwood JP, McCann GP, et al. Magnetic resonance perfusion or fractional flow reserve in coronary disease. *N Engl J Med*. 2019;380:2418-2428.

271. Knuuti J, Ballo H, Juarez-Orozco LE, et al. The performance of non-invasive tests to rule-in and rule-out significant coronary artery stenosis in patients with stable angina: a meta-analysis focused on post-test disease probability. *Eur Heart J*. 2018;39:3322-3330.

272. Danad I, Rajmakers PG, Driessen RS, et al. Comparison of coronary CT angiography, SPECT, PET, and hybrid imaging for diagnosis of ischemic heart disease determined by fractional flow reserve. *JAMA Cardiol*. 2017;2:1100-1107.

273. Neglia D, Rovai D, Caselli C, et al. Detection of significant coronary artery disease by noninvasive anatomical and functional imaging. *Circ Cardiovasc Imaging*. 2015;8:e002179.

274. Danad I, Szymonifka J, Twisk JWR, et al. Diagnostic performance of cardiac imaging methods to diagnose ischaemia-causing coronary artery disease when directly compared with fractional flow reserve as a reference standard: a meta-analysis. *Eur Heart J*. 2017;38:991-998.

275. Mieres JH, Gulati M, Bairey Merz N, et al. Role of noninvasive testing in the clinical evaluation of women with suspected ischemic heart disease: a consensus statement from the American Heart Association. *Circulation*. 2014;130:350-379.

276. Mark DB, Shaw L, Harrell FE Jr, et al. Prognostic value of a treadmill exercise score in outpatients with suspected coronary artery disease. *N Engl J Med*. 1991;325:849-853.

277. Lauer MS, Pothier CE, Magid DJ, et al. An externally validated model for predicting long-term survival after exercise treadmill testing in patients with suspected coronary artery disease and a normal electrocardiogram. *Ann Intern Med*. 2007;147:821-828.

278. Gulati M, Black HR, Shaw LJ, et al. The prognostic value of a nomogram for exercise capacity in women. *N Engl J Med*. 2005;353:468-475.

279. Ardestani A, Ahlberg AW, Katten DM, et al. Risk stratification using line source attenuation correction with rest/stress Tc-99m sestamibi SPECT myocardial perfusion imaging. *J Nucl Cardiol*. 2014;21:118-126.

280. van Dijk JD, Mouden M, Ottervanger JP, et al. Value of attenuation correction in stress-only myocardial perfusion imaging using CZT-SPECT. *J Nucl Cardiol*. 2017;24:395-401.

281. Gutstein A, Bental T, Solodky A, et al. Prognosis of stress-only SPECT myocardial perfusion imaging with prone imaging. *J Nucl Cardiol*. 2018;25:809-816.

282. Huang JY, Yen RF, Lee WC, et al. Improved diagnostic accuracy of thallium-201 myocardial perfusion single-photon emission computed tomography with CT attenuation correction. *J Nucl Cardiol*. 2019;26:1584-1595.

283. Huang JY, Huang CK, Yen RF, et al. Diagnostic performance of attenuation-corrected myocardial perfusion imaging for coronary artery disease: a systematic review and meta-analysis. *J Nucl Med*. 2016;57:1893-1898.

284. Ito S, Endo A, Okada T, et al. Comparison of CTAC and prone imaging for the detection of coronary artery disease using CZT SPECT. *Ann Nucl Med*. 2017;31:629-635.

285. Gibbons RJ, Carryer D, Liu H, et al. Use of echocardiography in outpatients with chest pain and normal resting electrocardiograms referred to Mayo Clinic Rochester. *Am Heart J*. 2018;196:49-55.

- 286.** Douglas PS, De Bruyne B, Pontone G, et al. 1-Year outcomes of FFRCT-guided care in patients with suspected coronary disease: the PLATFORM study. *J Am Coll Cardiol.* 2016;68:435-445.
- 287.** Min JK, Leipsic J, Pencina MJ, et al. Diagnostic accuracy of fractional flow reserve from anatomic CT angiography. *JAMA.* 2012;308:1237-1245.
- 288.** Andreini D, Modolo R, Katagiri Y, et al. Impact of fractional flow reserve derived from coronary computed tomography angiography on heart team treatment decision-making in patients with multivessel coronary artery disease: insights from the SYNTAX III REVOLUTION trial. *Circ Cardiovasc Interv.* 2019;12:e007607.
- 289.** Abidov A, Gallagher MJ, Chinnaiyan KM, et al. Clinical effectiveness of coronary computed tomographic angiography in the triage of patients to cardiac catheterization and revascularization after inconclusive stress testing: results of a 2-year prospective trial. *J Nucl Cardiol.* 2009;16:701-713.
- 290.** Christman MP, Bittencourt MS, Hulten E, et al. Yield of downstream tests after exercise treadmill testing: a prospective cohort study. *J Am Coll Cardiol.* 2014;63:1264-1274.
- 291.** Shaw LJ, Hausleiter J, Achenbach S, et al. Coronary computed tomographic angiography as a gatekeeper to invasive diagnostic and surgical procedures: results from the multicenter CONFIRM (Coronary CT Angiography Evaluation for Clinical Outcomes: an International Multicenter) registry. *J Am Coll Cardiol.* 2012;60:2103-2114.
- 292.** Schepis T, Gaemperli O, Koepfli P, et al. Added value of coronary artery calcium score as an adjunct to gated SPECT for the evaluation of coronary artery disease in an intermediate-risk population. *J Nucl Med.* 2007;48:1424-1430.
- 293.** Bavishi C, Argulian E, Chatterjee S, et al. CACS and the frequency of stress-induced myocardial ischemia during MPI: a meta-analysis. *J Am Coll Cardiol Img.* 2016;9:580-589.
- 294.** Rozanski A, Gransar H, Wong ND, et al. Use of coronary calcium scanning for predicting inducible myocardial ischemia: influence of patients' clinical presentation. *J Nucl Cardiol.* 2007;14:669-679.
- 295.** Brodov Y, Gransar H, Dey D, et al. Combined quantitative assessment of myocardial perfusion and coronary artery calcium score by hybrid 82Rb PET/CT improves detection of coronary artery disease. *J Nucl Med.* 2015;56:1345-1350.
- 296.** Ghadri JR, Pazhenkottil AP, Nkoulou RN, et al. Very high coronary calcium score unmasks obstructive coronary artery disease in patients with normal. *SPECT MPI. Heart.* 2011;97:998-1003.
- 297.** Schenker MP, Dorbala S, Hong EC, et al. Interrelation of coronary calcification, myocardial ischemia, and outcomes in patients with intermediate likelihood of coronary artery disease: a combined positron emission tomography/computed tomography study. *Circulation.* 2008;117:1693-1700.
- 298.** Marwick TH, Case C, Vasey C, et al. Prediction of mortality by exercise echocardiography: a strategy for combination with the Duke Treadmill Score. *Circulation.* 2001;103:2566-2571.
- 299.** Sicari R, Nihoyannopoulos P, Evangelista A, et al. Stress echocardiography expert consensus statement—executive summary: European Association of Echocardiography (EAE) (a registered branch of the ESC). *Eur Heart J.* 2009;30:278-289.
- 300.** Vitola JV, Wanderley MR Jr, Cerci RJ, et al. Outcome of patients with high-risk Duke Treadmill Score and normal myocardial perfusion imaging on SPECT. *J Nucl Cardiol.* 2016;23:1291-1300.
- 301.** Boiten HJ, van Domburg RT, Valkema R, et al. Eleven-year prognostic value of dobutamine stress (99m)Tc-sestamibi myocardial perfusion imaging in patients with limited exercise capacity. *Am J Cardiol.* 2015;115:884-889.
- 302.** Uretsky S, Supariwala A, Gurram S, et al. The interaction of exercise ability and body mass index upon long-term outcomes among patients undergoing stress-rest perfusion single-photon emission computed tomography imaging. *Am Heart J.* 2013;166:127-133.
- 303.** Bourque JM, Charlton GT, Holland BH, et al. Prognosis in patients achieving ≥ 10 METS on exercise stress testing: was SPECT imaging useful? *J Nucl Cardiol.* 2011;18:230-237.
- 304.** BARI 2D Study Group, Frye RL, August P, et al. A randomized trial of therapies for type 2 diabetes and coronary artery disease. *N Engl J Med.* 2009;360:2503-2515.
- 305.** Arnett DK, Blumenthal R, Albert M, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2019;74:e177-e232.
- 306.** Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2019;73:e285-e350.
- 307.** De Bruyne B, Pijls NH, Kalesan B, et al. Fractional flow reserve-guided PCI versus medical therapy in stable coronary disease. *N Engl J Med.* 2012;367:991-1001.
- 308.** De Bruyne B, Fearon WF, Pijls NH, et al. Fractional flow reserve-guided PCI for stable coronary artery disease. *N Engl J Med.* 2014;371:1208-1217.
- 309.** Tonino PA, De Bruyne B, Pijls NH, et al. Fractional flow reserve versus angiography for guiding percutaneous coronary intervention. *N Engl J Med.* 2009;360:213-224.
- 310.** Bhatt DL. Assessment of stable coronary lesions. *N Engl J Med.* 2017;376:1879-1881.
- 311.** Suh YJ, Hong YJ, Lee HJ, et al. Accuracy of CT for selecting candidates for coronary artery bypass graft surgery: combination with the SYNTAX score. *Radiology.* 2015;276:390-399.
- 312.** Chan M, Ridley L, Dunn DJ, et al. A systematic review and meta-analysis of multi-detector computed tomography in the assessment of coronary artery bypass grafts. *Int J Cardiol.* 2016;221:898-905.
- 313.** Small GR, Yam Y, Chen L, et al. Prognostic assessment of coronary artery bypass patients with 64-slice computed tomography angiography: anatomical information is incremental to clinical risk prediction. *J Am Coll Cardiol.* 2011;58:2389-2395.
- 314.** Collet C, Onuma Y, Andreini D, et al. Coronary computed tomography angiography for heart team decision-making in multivessel coronary artery disease. *Eur Heart J.* 2018;39:3689-3698.
- 315.** Shaw LJ, Berman DS, Maron DJ, et al. Optimal medical therapy with or without percutaneous coronary intervention to reduce ischemic burden: results from the Clinical Outcomes Utilizing Revascularization and Aggressive Drug Evaluation (COURAGE) trial nuclear substudy. *Circulation.* 2008;117:1283-1291.
- 316.** Shaw LJ, Cerqueira MD, Brooks MM, et al. Impact of left ventricular function and the extent of ischemia and scar by stress myocardial perfusion imaging on prognosis and therapeutic risk reduction in diabetic patients with coronary artery disease: results from the Bypass Angioplasty Revascularization Investigation 2 Diabetes (BARI 2D) trial. *J Nucl Cardiol.* 2012;19:658-669.
- 317.** Zellweger MJ, Fahrni G, Ritter M, et al. Prognostic value of "routine" cardiac stress imaging 5 years after percutaneous coronary intervention: the prospective long-term observational BASKET (Basel Stent Kosteffektivitats Trial) LATE IMAGING study. *J Am Coll Cardiol Intv.* 2014;7:615-621.
- 318.** Zellweger MJ, Kaiser C, Jeger R, et al. Coronary artery disease progression late after successful stent implantation. *J Am Coll Cardiol.* 2012;59:793-799.
- 319.** Zellweger MJ, Kaiser C, Brunner-La Rocca HP, et al. Value and limitations of target-vessel ischemia in predicting late clinical events after drug-eluting stent implantation. *J Nucl Med.* 2008;49:550-556.
- 320.** Shaw LJ, Weintraub WS, Maron DJ, et al. Baseline stress myocardial perfusion imaging results and outcomes in patients with stable ischemic heart disease randomized to optimal medical therapy with or without percutaneous coronary intervention. *Am Heart J.* 2012;164:243-250.
- 321.** Patel KK, Spertus JA, Arnold SV, et al. Ischemia on PET MPI may identify patients with improvement in angina and health status post-revascularization. *J Am Coll Cardiol.* 2019;74:1734-1736.
- 322.** Patel KK, Spertus JA, Chan PS, et al. Extent of myocardial ischemia on positron emission tomography and survival benefit with early revascularization. *J Am Coll Cardiol.* 2019;74:1645-1654.
- 323.** Reynolds HR, Shaw LJ, Min JK, et al. Association of sex with severity of coronary artery disease, ischemia, and symptom burden in patients with moderate or severe ischemia: secondary analysis of the ISCHEMIA randomized clinical trial. *JAMA Cardiol.* 2020;5:1-14.
- 324.** Schwitzer J, Wacker CM, van Rossum AC, et al. MR-IMPACT: comparison of perfusion-cardiac magnetic resonance with single-photon emission computed tomography for the detection of coronary artery disease in a multicentre, multivendor, randomized trial. *Eur Heart J.* 2008;29:480-489.
- 325.** Arai AE, Schulz-Menger J, Berman D, et al. Gadobutrol-enhanced cardiac magnetic resonance imaging for detection of coronary artery disease. *J Am Coll Cardiol.* 2020;76:1536-1547.
- 326.** Takx RA, Blomberg BA, El Aidi H, et al. Diagnostic accuracy of stress myocardial perfusion imaging compared to invasive coronary angiography with

- fractional flow reserve meta-analysis. *Circ Cardiovasc Imaging*. 2015;8.
- 327.** Heitner JF, Kim RJ, Kim HW, et al. Prognostic value of vasodilator stress cardiac magnetic resonance imaging: a multicenter study with 48000 patient-years of follow-up. *JAMA Cardiol*. 2019;4:256-264.
- 328.** Kato S, Saito N, Nakachi T, et al. Stress perfusion coronary flow reserve versus cardiac magnetic resonance for known or suspected CAD. *J Am Coll Cardiol*. 2017;70:869-879.
- 329.** Vincenti G, Masci PG, Monney P, et al. Stress perfusion CMR in patients with known and suspected CAD: prognostic value and optimal ischemic threshold for revascularization. *J Am Coll Cardiol Img*. 2017;10:526-537.
- 330.** Kwong RY, Ge Y, Steel K, et al. Cardiac magnetic resonance stress perfusion imaging for evaluation of patients with chest pain. *J Am Coll Cardiol*. 2019;74:1741-1755.
- 331.** Taqueti VR, Hachamovitch R, Murthy VL, et al. Global coronary flow reserve is associated with adverse cardiovascular events independently of luminal angiographic severity and modifies the effect of early revascularization. *Circulation*. 2015;131:19-27.
- 332.** Driessen RS, Danad I, Stuijzand WJ, et al. Comparison of coronary computed tomography angiography, fractional flow reserve, and perfusion imaging for ischemia diagnosis. *J Am Coll Cardiol*. 2019;73:161-173.
- 333.** Patel KK, Spertus JA, Chan PS, et al. Myocardial blood flow reserve assessed by positron emission tomography myocardial perfusion imaging identifies patients with a survival benefit from early revascularization. *Eur Heart J*. 2020;41:759-768.
- 334.** Bom MJ, van Diemen PA, Driessen RS, et al. Prognostic value of [15O]H₂O positron emission tomography-derived global and regional myocardial perfusion. *Eur Heart J Cardiovasc Imaging*. 2020;21:777-786.
- 335.** Knott KD, Seraphim A, Augusto JB, et al. The prognostic significance of quantitative myocardial perfusion: an artificial intelligence-based approach using perfusion mapping. *Circulation*. 2020;141:1282-1291.
- 336.** Ho PM, Rumsfeld JS, Peterson PN, et al. Chest pain on exercise treadmill test predicts future cardiac hospitalizations. *Clin Cardiol*. 2007;30:505-510.
- 337.** Hambrecht R, Walther C, Mobius-Winkler S, et al. Percutaneous coronary angioplasty compared with exercise training in patients with stable coronary artery disease: a randomized trial. *Circulation*. 2004;109:1371-1378.
- 338.** Jaureguizar KV, Vicente-Campos D, Bautista LR, et al. Effect of high-intensity interval versus continuous exercise training on functional capacity and quality of life in patients with coronary artery disease: a randomized clinical trial. *J Cardiopulm Rehabil Prev*. 2016;36:96-105.
- 339.** Barbero U, Iannaccone M, d'Ascenzo F, et al. 64 slice-coronary computed tomography sensitivity and specificity in the evaluation of coronary artery bypass graft stenosis: a meta-analysis. *Int J Cardiol*. 2016;216:52-57.
- 340.** Lee SE, Chang HJ, Sung JM, et al. Effects of statins on coronary atherosclerotic plaques: the PARADIGM study. *J Am Coll Cardiol Img*. 2018;11:1475-1484.
- 341.** Lee SE, Sung JM, Rizvi A, et al. Quantification of coronary atherosclerosis in the assessment of coronary artery disease. *Circ Cardiovasc Imaging*. 2018;11:e007562.
- 342.** Williams MC, Kwiecinski J, Doris M, et al. Low-attenuation noncalcified plaque on coronary computed tomography angiography predicts myocardial infarction: results from the Multicenter SCOT-HEART Trial (Scottish Computed Tomography of the HEART). *Circulation*. 2020;141:1452-1462.
- 343.** Ferencik M, Mayrhofer T, Bittner DO, et al. Use of high-risk coronary atherosclerotic plaque detection for risk stratification of patients with stable chest pain: a secondary analysis of the PROMISE randomized clinical trial. *JAMA Cardiol*. 2018;3:144-152.
- 344.** Taqueti VR, Solomon SD, Shah AM, et al. Coronary microvascular dysfunction and future risk of heart failure with preserved ejection fraction. *Eur Heart J*. 2018;39:840-849.
- 345.** Taqueti VR, Shaw LJ, Cook NR, et al. Excess cardiovascular risk in women relative to men referred for coronary angiography is associated with severely impaired coronary flow reserve, not obstructive disease. *Circulation*. 2017;135:566-577.
- 346.** Indorkar R, Kwong RY, Romano S, et al. Global coronary flow reserve measured during stress cardiac magnetic resonance imaging is an independent predictor of adverse cardiovascular events. *J Am Coll Cardiol Img*. 2019;12:1686-1695.
- 347.** Zorach B, Shaw PW, Bourque J, et al. Quantitative cardiovascular magnetic resonance perfusion imaging identifies reduced flow reserve in microvascular coronary artery disease. *J Cardiovasc Magn Reson*. 2018;20:14.
- 348.** Bairey Merz CN, Pepine CJ, Walsh MN, et al. Ischemia and No Obstructive Coronary Artery Disease (INOCA): developing evidence-based therapies and research agenda for the next decade. *Circulation*. 2017;135:1075-1092.
- 349.** AlBadri A, Bairy Merz CN, Johnson BD, et al. Impact of abnormal coronary reactivity on long-term clinical outcomes in women. *J Am Coll Cardiol*. 2019;73:684-693.
- 350.** AlBadri A, Sharif B, Wei J, et al. Intracoronary bolus injection versus intravenous infusion of adenosine for assessment of coronary flow velocity reserve in women with signs and symptoms of myocardial ischemia and no obstructive coronary artery disease. *J Am Coll Cardiol Intv*. 2018;11:2125-2127.
- 351.** Ford TJ, Stanley B, Sidik N, et al. 1-Year Outcomes of Angina Management Guided by Invasive Coronary Function Testing (CorMicA). *J Am Coll Cardiol Intv*. 2020;13:33-45.
- 352.** Ford TJ, Corcoran D, Sidik N, et al. Coronary microvascular dysfunction: assessment of both structure and function. *J Am Coll Cardiol*. 2018;72:584-586.

KEY WORDS ACC/AHA Clinical Practice Guidelines, chest pain, angina, coronary artery disease, acute coronary syndrome, myocardial ischemia, myocardial infarction, myocardial injury, noncardiac, accelerated diagnostic pathway, clinical decision pathway, sex differences, troponins, chest pain syndromes, biomarkers, shared-decision making, noncardiac chest pain, cardiac imaging