

Search...



[OVERVIEW](#) [**POCKET GUIDE**](#) [FULL TEXT LINK](#)



Evaluation and Diagnosis of Chest Pain

Last updated May 24, 2022

Key Points



Scope of the Problem

Synopsis

- After injuries, chest pain is the second most common reason for adults to present to the emergency department (ED) in the United States and accounts for >6.5 million visits, which is 4.7% of all ED visits.
- Chest pain also leads to nearly 4 million outpatient visits annually in the United States.
- Chest pain remains a diagnostic challenge in the ED and outpatient setting and requires thorough clinical evaluation.
 - Although the cause of chest pain is often noncardiac, coronary artery disease (CAD) affects >18.2 million adults in the United States and remains the leading cause of death for men and women, accounting for >365,000 deaths

FEEDBACK

- Distinguishing between serious and benign causes of chest pain is imperative.
- The lifetime prevalence of chest pain in the United States is 20% to 40%, and women experience this symptom more often than men.
- Of all ED patients with chest pain, only 5.1% will have an acute coronary syndrome (ACS), and more than half will ultimately be found to have a noncardiac cause.
- Nonetheless, chest pain is the most common symptom of CAD in both men and women.

Top 10 Take-Home Messages:*

- 1. Chest Pain Means More Than Pain in the Chest.** Pain, pressure, tightness, or discomfort in the chest, shoulders, arms, neck, back, upper abdomen, or jaw, as well as shortness of breath and fatigue should all be considered anginal equivalents.
- 2. High-Sensitivity Troponins Preferred.** High-sensitivity cardiac troponins are the preferred standard for establishing a biomarker diagnosis of acute myocardial infarction, allowing for more accurate detection and exclusion of myocardial injury.
- 3. Early Care for Acute Symptoms.** Patients with acute chest pain or chest pain equivalent symptoms should seek medical care immediately by calling 9-1-1. Although most patients will not have a cardiac cause, the evaluation of all patients should focus on the early identification or exclusion of life-threatening causes.
- 4. Share the Decision-Making.** Clinically stable patients presenting with chest pain should be included in decision-making; information about risk of adverse events, radiation exposure, costs, and alternative options should be provided to facilitate the discussion.
- 5. Testing Not Needed Routinely for Low-Risk Patients.** For patients with acute or stable chest pain determined to be low risk, urgent diagnostic testing for suspected coronary artery disease is not needed.
- 6. Pathways.** Clinical decision pathways for chest pain in the emergency department and outpatient settings should be used routinely.
- 7. Accompanying Symptoms.** Chest pain is the dominant and most frequent symptom for both men and women ultimately diagnosed with Acute Coronary Syndrome. Women may be more likely to present with accompanying symptoms such as nausea and shortness of breath.
- 8. Identify Patients Most Likely to Benefit From Further Testing.** Patients with acute or stable chest pain who are at intermediate risk or intermediate to high pre-test risk of obstructive coronary artery disease, respectively, will benefit the most from cardiac imaging and testing.
- 9. Noncardiac Is In. Atypical Is Out.** “Noncardiac” should be used if heart disease is not suspected. “Atypical” is a misleading descriptor of chest pain, and its use is discouraged.
- 10. Structured Risk Assessment Should Be Used.** For patients presenting with acute or stable chest pain, risk for coronary artery disease and adverse events should be estimated using evidence-based diagnostic protocols.

* Figure 1 illustrates the take-home messages.

Figure 1. Take-Home Messages for the Evaluation and Diagnosis of Chest Pain




Figure 1. Take-Home Messages for the Evaluation and Diagnosis of Chest Pain Cont'd



Treatment

1.4.2. Defining Chest Pain

- ▶ 1. An initial assessment of chest pain is recommended to triage patients effectively on the basis of the likelihood that symptoms may be attributable to myocardial ischemia. (*I, B-NR*)
- ▶ 2. Chest pain should not be described as atypical, because it is not helpful in determining the cause and can be misinterpreted as benign in nature. Instead, chest pain should be described as cardiac, possibly cardiac, or noncardiac because these terms are more specific to the potential underlying diagnosis. (*I, C-LD*)

2. Initial Evaluation

2.1 Recommendation for History



1. In patients with chest pain, a focused history that includes characteristics and duration of symptoms relative to presentation as well as associated features, and cardiovascular risk factor assessment should be obtained.. (I, C-LD)

Figure 2. Index of Suspicion That Chest "Pain" Is Ischemic in Origin on the Basis of Commonly Used Descriptors






Figure 3. Top 10 Causes of Chest Pain in the ED Based on Age (Weighted Percentage)




- Painful respiration
- Abdominal pain
- Bone/musculoskeletal
- Anxiety
- Superficial contusion
- Cardiac dysrhythmia
- Esophageal disorder
- Other upper respiratory infection
- Other & unspecified lower respiratory infection

- Coronary atherosclerosis
- Painful respiration
- Acute myocardial infarction
- Cardiac dysrhythmia
- Abdominal pain
- Pneumonia
- Esophageal disorder
- Superficial injury; contusion
- Essential hypertension

65-79 y of Age



≥80 y of Age



- Nonspecific chest pain
- Coronary atherosclerosis
- Cardiac dysrhythmia
- Acute myocardial infarction
- Painful respiration
- Congestive heart failure
- Abdominal pain
- Pneumonia
- Other & unspecified lower respiratory disease

- Nonspecific chest pain
- Coronary atherosclerosis
- Congestive heart failure
- Acute myocardial infarction
- Pneumonia
- Painful respiration
- Cardiac dysrhythmia
- Abdominal pain
- Other & unspecified lower respiratory disease

Created using data from Hsia RY, et al. *Intern Med*.2016;176:1029-32.

Note: The numbering of the following tables and figures differs from that of the Clinical Practice Guideline.

Table 1. Chest Pain Characteristics and Corresponding Causes

<p>Nature</p> <ul style="list-style-type: none"> • Anginal symptoms are perceived as retrosternal chest discomfort (e.g., pain, discomfort, heaviness, tightness, pressure, constriction, squeezing) (Section 1.4.2, Defining Chest Pain). • Sharp chest pain that increases with inspiration and lying supine is unlikely related to ischemic heart disease (e.g., these symptoms usually occur with acute pericarditis).
<p>Onset and duration</p> <ul style="list-style-type: none"> • Anginal symptoms gradually build in intensity over a few minutes. • Sudden onset of ripping chest pain (with radiation to the upper or lower back) is unlikely to be anginal and is suspicious of an acute aortic syndrome. • Fleeting chest pain—of few seconds' duration—is unlikely to be related to ischemic heart disease.
<p>Location and radiation</p> <ul style="list-style-type: none"> • Pain that can be localized to a very limited area and pain radiating to below the umbilicus or hip are unlikely related to myocardial ischemia.
<p>Severity</p>

2.1.1. Focus on the Uniqueness of Chest Pain in Women

- ▶
 1. Women who present with chest pain are at risk for underdiagnosis, and potential cardiac causes should always be considered. *(I, B-NR)*
- ▶
 2. In women presenting with chest pain, it is recommended to obtain a history that emphasizes accompanying symptoms that are more common in women with ACS. *(I, B-NR)*

2.1.2. Considerations for Older Patients With Chest Pain

- ▶
 1. In patients with chest pain who are >75 years of age, ACS should be considered when accompanying symptoms such as shortness of breath, syncope, or acute delirium are present, or when an unexplained fall has occurred. *(I, C-LD)*

2.1.3. Considerations for Diverse Patient Populations With Chest Pain

- ▶
 1. Cultural competency training is recommended to help achieve the best outcomes in patients of diverse racial and ethnic backgrounds who present with chest pain. *(I, C-LD)*
- ▶
 2. Among patients of diverse race and ethnicity presenting with chest pain in whom English may not be their primary language, addressing language barriers with the use of formal translation services is recommended. *(I, C-LD)*

2.1.4. Patient-Centric Considerations

- ▶
 1. In patients with acute chest pain, it is recommended that 9-1-1 be activated by patients or bystanders to initiate transport to the closest ED by emergency medical services (EMS). *(I, C-LD)*

2.2. Physical Examination



1. In patients presenting with chest pain, a focused cardiovascular examination should be performed initially to aid in the diagnosis of ACS or other potentially serious causes of chest pain (e.g., aortic dissection, PE, or esophageal rupture) and to identify complications. (*I, C-EO*)

TABLE 2. PHYSICAL EXAMINATION IN PATIENTS WITH CHEST PAIN

Clinical Syndrome	Findings
Emergency	
ACS	<ul style="list-style-type: none">• Diaphoresis, tachypnea, tachycardia, hypotension, crackles, S3, MR murmur; examination may be normal in uncomplicated cases
PE	<ul style="list-style-type: none">• Tachycardia + dyspnea—>90% of patients; pain with inspiration
Aortic dissection	<ul style="list-style-type: none">• Connective tissue disorders (e.g., Marfan syndrome), extremity pulse differential (30% of patients, type A>B)• Severe pain, abrupt onset + pulse differential + widened mediastinum on CXR >80% probability of dissection• Frequency of syncope >10%, AR 40%–75% (type A)
Esophageal rupture	<ul style="list-style-type: none">• Emesis, subcutaneous emphysema, pneumothorax (20% patients), unilateral decreased or absent breath sounds
Other	
Noncoronary cardiac: AS, AR,	<ul style="list-style-type: none">• AS: Characteristic systolic murmur, tardus or parvus carotid pulse

2.3. Diagnostic Testing

2.3.1. Setting Considerations



1. Unless a noncardiac cause is evident, an ECG should be performed for patients seen in the office setting with stable chest pain. If an ECG is unavailable the patient should be referred to the ED so one can be obtained. (*I, B-NR*)



2. Patients with clinical evidence of ACS or other life-threatening causes of acute chest pain seen in the office setting should be transported urgently to the ED, ideally by EMS. (*I, C-LD*)



3. In all patients who present with acute chest pain regardless of the setting, an ECG should be acquired and reviewed for STEMI within 10 minutes of arrival. (*I, C-LD*)



4. In all patients presenting to the ED with acute chest pain and suspected ACS, cTn should be measured as soon as possible after presentation. (*I, C-LD*)



5. For patients with acute chest pain and suspected ACS initially evaluated in the office setting, delayed transfer to the ED for cTn or other diagnostic testing should be avoided. (*III - Harm, C-LD*)


2.3.2. Electrocardiogram (ECG)



1. In patients with chest pain in which an initial ECG is nondiagnostic, serial ECGs to detect potential ischemic changes should be performed, especially when clinical suspicion of ACS is high, symptoms are persistent, or the clinical condition deteriorates. (*I, C-EO*)

- ▶
- 2. Patients with chest pain in whom the initial ECG is consistent with an ACS should be treated according to STEMI and NSTEMI-ACS guidelines. (*I, C-EO*)
- ▶
- 3. In patients with chest pain and intermediate-to-high clinical suspicion for ACS in whom the initial ECG is nondiagnostic, supplemental electrocardiographic leads V7 to V9 are reasonable to rule out posterior MI. (*IIa, B-NR*)

Figure 4. Electrocardiographic-Directed Management of Chest Pain




2.3.3. Chest Radiography

- ▶
- 1. In patients presenting with acute chest pain, a chest radiograph is useful to evaluate for other potential cardiac, pulmonary, and thoracic causes of symptoms. (*I, C-EO*)

2.3.4. Biomarkers

- ▶
- 1. In patients presenting with acute chest pain, serial cTn I or T levels are useful to identify abnormal values and a rising or falling pattern indicative of acute myocardial injury. (*I, B-NR*)
- ▶
- 2. In patients presenting with acute chest pain, high-sensitivity cTn is the preferred biomarker because it enables more rapid detection or exclusion of myocardial injury and increases diagnostic accuracy. (*I, B-NR*)
- ▶
- 3. Clinicians should be familiar with the analytical performance and the 99th percentile upper reference limit that defines myocardial injury for the cTn assay used at their institution. (*I, C-EO*)
- ▶
- 4. With availability of cTn, creatine kinase myocardial (CK-MB) isoenzyme and myoglobin are not useful for diagnosis of acute myocardial injury. (*III - No Benefit, B-NR*)

Figure 5. Chest Pain and Cardiac Testing Considerations



The choice of imaging depends on the clinical question of importance, to either a) ascertain the diagnosis of CAD and define coronary anatomy or b) assess ischemia severity among patients with an expected higher likelihood of ischemia with an abnormal resting ECG or those incapable of performing maximal exercise.

Please refer to Section 4.1.

For risk assessment in acute chest pain: See Figure 9.

For risk assessment in stable chest pain: See Figure 11.

Figure 6. Choosing the Right Diagnostic Test

Pretest likelihood of CAD	Low	→	No testing necessary	→	Option for CAC for ASCVD risk stratification
	Intermediate-high	→	Younger patient (<65 y of age)	OR	Less obstructive CAD suspected → CCTA favored
	Intermediate-high	→	Older patient (≥65 y of age)	OR	More obstructive CAD suspected → Stress testing favored

	Favors use of CCTA	Favors use of stress imaging
Goal	<ul style="list-style-type: none"> Rule out obstructive CAD Detect nonobstructive CAD 	<ul style="list-style-type: none"> Ischemia-guided management
Availability and expertise	<ul style="list-style-type: none"> High-quality imaging and expert interpretation routinely available 	<ul style="list-style-type: none"> High-quality imaging and expert interpretation routinely available
Likelihood of obstructive CAD	<ul style="list-style-type: none"> Age <65 y 	<ul style="list-style-type: none"> Age ≥65 y
Prior test results	<ul style="list-style-type: none"> Prior functional study inconclusive 	<ul style="list-style-type: none"> Prior CCTA inconclusive
Other compelling indications	<ul style="list-style-type: none"> Anomalous coronary arteries Require evaluation of aorta or pulmonary arteries 	<ul style="list-style-type: none"> Suspect scar (especially if PET or stress CMR available) Suspect coronary microvascular dysfunction (when PET or CMR available)

Stress testing information					
	ETT	Stress echocardiography	SPECT MPI	PET MPI	Stress CMR MPI
Patient capable of exercise	✓	✓	✓		
Pharmacologic stress indicated		✓	✓	✓	✓
Quantitative flow				✓	✓
LV dysfunction/scar		✓	✓	✓	

Table 3. Contraindication by Type of Imaging Modality and Stress Protocol


aExercise ECG	Stress Nuclear ^a	
<ul style="list-style-type: none"> • Abnormal ST changes on resting ECG, digoxin, left bundle branch block, Wolff-Parkinson-White pattern, ventricular paced rhythm (unless test is performed to establish exercise capacity and not for diagnosis of ischemia) • Unable to achieve ≥ 5 METs or unsafe to exercise • High-risk unstable angina or AMI (<2 d) i.e., active ACS • Uncontrolled heart failure • Significant cardiac arrhythmias (e.g., VT, complete atrioventricular block) or high risk for arrhythmias caused by QT prolongation • Severe symptomatic aortic stenosis • Severe systemic arterial hypertension (e.g., $\geq 200/110$ mm Hg) • Acute illness (e.g., acute PE, acute myocarditis/pericarditis, acute aortic dissection) 	<ul style="list-style-type: none"> • High-risk unstable angina, complicated ACS or AMI (<2 d) • Contraindications to vasodilator administration <ul style="list-style-type: none"> ◦ Significant arrhythmias (e.g., VT, second- or third-degree atrioventricular block) or sinus bradycardia <45 bpm ◦ Significant hypotension (SBP <90 mm Hg) ◦ Known or suspected bronchoconstrictive or bronchospastic disease ◦ Recent use of dipyridamole or dipyridamole-containing medications ◦ Use of methylxanthines (e.g., aminophylline, caffeine) within 12 hours ◦ Known hypersensitivity to adenosine, regadenoson • Severe systemic arterial hypertension (e.g., $\geq 200/110$ mm Hg) 	
Stress Echocardiography	Stress CMR	CCTA ^a


For all the imaging modalities, inability to achieve high-quality images should be considered, in particular for obese patients.

^a Screening for potential pregnancy by history and/or pregnancy testing should be performed according to the local imaging facilities policies for undertaking radiological examinations that involve ionizing radiation in women of child-bearing age.

^b Low-dose dobutamine may be useful for assessing for low-gradient AS.

Figure 7. Patient-Centric Algorithms for Acute Chest Pain





4.1 Patients With Acute Chest Pain and Suspected ACS (Not Including STEMI)

- ▶
- 1. In patients presenting with acute chest pain and suspected ACS, clinical decision pathways (CDPs) should categorize patients into low-, intermediate-, and high-risk strata to facilitate disposition and subsequent diagnostic evaluation. (I, B-NR)
- ▶

2. In the evaluation of patients presenting with acute chest pain and suspected ACS for whom serial troponins are indicated to exclude myocardial injury, recommended time intervals after the initial troponin sample collection (time zero) for repeat measurements are: 1 to 3 hours for high-sensitivity troponin and 3 to 6 hours for conventional troponin assays. **(I, B-NR)**
- ▶
3. To standardize the detection and differentiation of myocardial injury in patients presenting with acute chest pain and suspected ACS, institutions should implement a CDP that includes a protocol for troponin sampling based on their particular assay. **(I, C-LD)**
- ▶
4. In patients with acute chest pain and suspected ACS, previous testing when available should be considered and incorporated into CDPs. **(I, C-LD)**
- ▶
5. For patients with acute chest pain, a normal ECG, and symptoms suggestive of ACS that began at least 3 hours before ED arrival, a single hs-cTn concentration that is below the limit of detection on initial measurement (time zero) is reasonable to exclude myocardial injury. **(IIa, B-NR)**

Figure 8. General Approach to Risk Stratification of Patients With Suspected ACS




Table 4. Warranty Period for Prior Cardiac Testing

Test Modality	Result	Warranty Period
---------------	--------	-----------------

Test Modality	Result	Warranty Period
Anatomic	<ul style="list-style-type: none"> • Normal coronary angiogram • CCTA with no stenosis or plaque 	2 y
Stress testing	<ul style="list-style-type: none"> • Normal stress test (given adequate stress) 	1 y

4.1.1. Low-Risk Patients With Acute Chest Pain

- ▶
- 1. Patients with acute chest pain and a 30-day risk of death or MACE <1% should be designated as low risk. **(I, B-NR)**
- ▶
- 2. In patients with acute chest pain and suspected ACS who are deemed low-risk (<1% 30-day risk of death or MACE), it is reasonable to discharge home without admission or urgent cardiac testing. **(IIa, B-R)**

TABLE 5. DEFINITION USED FOR LOW-RISK PATIENTS WITH CHEST PAIN

Low Risk (<1% 30-d Risk for Death or MACE)	
hs-cTn Based	
T-0	T-0 hs-cTn below the assay limit of detection or “very low” threshold if symptoms present for at least 3 h
T-0 and 1- or 2-h Delta	T-0 hs-cTn and 1- or 2-h delta are both below the assay “low” thresholds (>99% NPV for 30-d MACE)
Clinical Decision Pathway Based	
HEART Pathway	HEART score <3, initial and serial cTn/hs-cTn < assay 99th percentile
EDACS	EDACS <16; initial and serial cTn/hs-cTn < assay 99th percentile
ADAPT	TIMI score 0, initial and serial cTn/hs-cTn < assay 99th percentile
mADAPT	TIMI score 0/1, initial and serial cTn/hs-cTn < assay 99th percentile
NOTR	0 factors

4.1.2. Intermediate-Risk Patients With Acute Chest Pain

- ▶
- 1. For intermediate-risk patients with acute chest pain, TTE is recommended as a rapid, bedside test to establish baseline ventricular and valvular function, evaluate for wall motion abnormalities, and to assess for pericardial effusion. **(I, C-EO)**
- ▶
- 2. For intermediate-risk patients with acute chest pain, management in an observation unit is reasonable to shorten length of stay and lower cost relative to an inpatient admission. **(IIa, A)**

4.1.2.1. Intermediate-Risk Patients With No Known (CAD)

ANATOMIC TESTING

- ▶
- 1. For intermediate-risk patients with acute chest pain and no known CAD eligible for diagnostic testing after a negative or inconclusive evaluation for ACS, CCTA is useful for exclusion of atherosclerotic plaque and obstructive CAD. **(I, A)**
- ▶
- 2. For intermediate-risk patients with acute chest pain, moderate-severe ischemia on current or prior (≤1 year) stress testing, and no known CAD established by prior anatomic testing, ICA is recommended. **(I, C-EO)**
- ▶

- For intermediate-risk patients with acute chest pain with evidence of previous mildly abnormal stress test results (≤ 1 year), CCTA is reasonable for diagnosing obstructive CAD. (*Ila, C-LD*)

STRESS TESTING



- For intermediate-risk patients with acute chest pain and no known CAD who are eligible for cardiac testing, either exercise ECG, stress echocardiography, stress PET/SPECT MPI, or stress CMR is useful for the diagnosis of myocardial ischemia. (*I, B-NR*)

SEQUENTIAL OR ADD-ON DIAGNOSTIC TESTING



- For intermediate-risk patients with acute chest pain and no known CAD, with a coronary artery stenosis of 40% to 90% in a proximal or middle coronary artery on CCTA, FFR-CT can be useful for the diagnosis of vessel-specific ischemia and to guide decision-making regarding the use of coronary revascularization. (*Ila, B-NR*)




- For intermediate-risk patients with acute chest pain and no known CAD, as well as an inconclusive prior stress test, CCTA can be useful for excluding the presence of atherosclerotic plaque and obstructive CAD. (*Ila, C-EO*)



- For intermediate-risk patients with acute chest pain and no known CAD, with an inconclusive CCTA, stress imaging (with echocardiography, PET/SPECT MPI, or CMR) can be useful for the diagnosis of myocardial ischemia. (*Ila, C-EO*)

Figure 9. Evaluation Algorithm for Patients With Suspected ACS at Intermediate Risk With No Known CAD



4.1.2.2. Intermediate-Risk Patients With Acute Chest Pain and Known CAD




- For intermediate-risk patients with acute chest pain who have known CAD and present with new onset or worsening symptoms, GDMT should be optimized before additional cardiac testing is performed. (*I, A*)



2. For intermediate-risk patients with acute chest pain who have worsening frequency of symptoms with significant left main, proximal left anterior descending stenosis, or multivessel CAD on prior anatomic testing or history of prior coronary revascularization, ICA is recommended. (*I, A*)
- ▶
3. For intermediate-risk patients with acute chest pain and known nonobstructive CAD, CCTA can be useful to determine progression of atherosclerotic plaque and obstructive CAD. (*IIa, B-NR*)
- ▶
4. For intermediate-risk patients with acute chest pain and coronary artery stenosis of 40% to 90% in a proximal or middle segment on CCTA, FFR-CT is reasonable for diagnosis of vessel-specific ischemia and to guide decision-making regarding the use of coronary revascularization. (*IIa, B-NR*)
- ▶
5. For intermediate-risk patients with acute chest pain and known CAD who have new onset or worsening symptoms, stress imaging (PET/SPECT MPI, CMR, or stress echocardiography) is reasonable. (*IIa, B-NR*)

Figure 10. Evaluation Algorithm for Patients With Suspected ACS at Intermediate Risk With Known CAD



4.1.3. High-Risk Patients With Acute Chest Pain

- ▶
1. For patients with acute chest pain and suspected ACS who have new ischemic changes on electrocardiography, troponin-confirmed acute myocardial injury, new-onset left ventricular systolic dysfunction (ejection fraction <40%), newly diagnosed moderate-severe ischemia on stress testing, hemodynamic instability, and/or a high clinical decision pathway (CDP) risk score should be designated as high risk for short-term MACE. (*I, B-NR*)
- ▶
2. For patients with acute chest pain and suspected ACS who are designated as high risk, ICA is recommended. (*I, C-EO*)
- ▶
3. For high-risk patients with acute chest pain who are troponin positive in whom obstructive CAD has been excluded by CCTA or ICA, CMR or echocardiography can be effective in establishing alternative diagnoses. (*IIa, B-NR*)

4.1.4. Acute Chest Pain in Patients With Prior CABG Surgery

- ▶
 1. In patients with prior CABG surgery presenting with acute chest pain who do not have ACS, performing stress imaging is effective to evaluate for myocardial ischemia or CCTA for graft stenosis or occlusion. (*I, C-LD*)
- ▶
 2. In patients with prior CABG surgery presenting with acute chest pain, who do not have ACS or who have an indeterminate/nondiagnostic stress test, ICA is useful. (*I, C-LD*)

4.1.5. Evaluation of Patients With Acute Chest Pain Receiving Dialysis

- ▶
 1. In patients who experience acute unremitting chest pain while undergoing dialysis, transfer by EMS to an acute care setting is recommended. (*I, B-NR*)

4.1.6. Evaluation of Acute Chest Pain in Patients With Cocaine and Methamphetamine Use

- ▶
 1. In patients presenting with acute chest pain, it is reasonable to consider cocaine and methamphetamine use as a cause of their symptoms. (*IIa, B-NR*)

4.1.7. Shared Decision-Making in Patients With Acute Chest Pain

- ▶
 1. For patients with acute chest pain and suspected ACS who are deemed low risk by a CDP, patient decision aids are beneficial to improve understanding and effectively facilitate risk communication. (*I, B-R*)
- ▶
 2. For patients with acute chest pain and suspected ACS who are deemed intermediate risk by a CDP, shared decision-making between the clinician and patient regarding the need for admission, for observation, discharge, or further evaluation in an outpatient setting is recommended for improving patient understanding and reducing low-value testing. (*I, B-R*)

4.2. Evaluation of Acute Chest Pain With Nonischemic Cardiac Pathologies

- ▶
 1. In patients with acute chest pain in whom other potentially life-threatening nonischemic cardiac conditions are suspected (e.g., aortic pathology, pericardial effusion, endocarditis), TTE is recommended for diagnosis. (*I, C-EO*)

4.2.1. Acute Chest Pain With Suspected Acute Aortic Syndrome

- ▶
 1. In patients with acute chest pain where there is clinical concern for aortic dissection, computed tomography angiography (CTA) of the chest, abdomen, and pelvis is recommended for diagnosis and treatment planning. (*I, C-EO*)
- ▶
 2. In patients with acute chest pain where there is clinical concern for aortic dissection, TEE or CMR should be performed to make the diagnosis if CT is contraindicated or unavailable. (*I, C-EO*)

4.2.2. Acute Chest Pain With Suspected PE

- ▶
 1. In stable patients with acute chest pain with high clinical suspicion for PE, CTA using a PE protocol is recommended. (*I, B-NR*)
- ▶
 2. For patients with acute chest pain and possible PE, need for further testing should be guided by pretest probability. (*I, C-EO*)

4.2.3. Acute Chest Pain With Suspected Myopericarditis

- ▶
 1. In patients with acute chest pain and myocardial injury who have nonobstructive coronary arteries on anatomic testing, CMR with gadolinium contrast is effective to distinguish myopericarditis from other causes, including myocardial

and nonobstructive coronary arteries (MINOCA). (*I, B-NR*)

- ▶
- 2. In patients with acute chest pain with suspected acute myopericarditis, CMR is useful if there is diagnostic uncertainty, or to determine the presence and extent of myocardial and pericardial inflammation and fibrosis. (*I, B-NR*)
- ▶
- 3. In patients with acute chest pain and suspected myopericarditis, TTE is effective to determine the presence of ventricular wall motion abnormalities, pericardial effusion, valvular abnormalities, or restrictive physiology. (*I, C-EO*)
- ▶
- 4. In patients with acute chest pain with suspected acute pericarditis, non-contrast or contrast cardiac CT scanning may be reasonable to determine the presence and degree of pericardial thickening. (*I**b**, C-LD*)

4.2.4. Acute Chest Pain With Valvular Heart Disease

- ▶
- 1. In patients presenting with acute chest pain with suspected or known history of valvular heart disease (VHD), TTE is useful in determining the presence, severity, and cause of VHD. (*I, C-EO*)
- ▶
- 2. In patients presenting with acute chest pain with suspected or known VHD in whom TTE diagnostic quality is inadequate, TEE (with 3D imaging if available) is useful in determining the severity and cause of VHD. (*I, C-EO*)
- ▶
- 3. In patients presenting with acute chest pain with known or suspected VHD, CMR imaging is reasonable as an alternative to TTE and/or TEE is nondiagnostic. (*I**a**, C-EO*)

4.3 Evaluation of Acute Chest Pain With Suspected Noncardiac Causes

- ▶
- 1. Patients with acute chest pain should be evaluated for noncardiac causes if they have persistent or recurring symptoms despite a negative stress test or anatomic cardiac evaluation, or a low-risk designation by a CDP. (*I, C-EO*)

Table 6. Differential Diagnosis of Noncardiac Chest Pain

Respiratory	Pulmonary embolism	Bronchitis
	Pneumothorax/hemothorax	Pleural irritation
	Pneumomediastinum	Malignancy
	Pneumonia	
Gastrointestinal	Cholecystitis	Peptic ulcer disease
	Pancreatitis	Esophageal spasm
	Hiatal hernia	Dyspepsia
	Gastroesophageal reflux disease/gastritis/esophagitis	
Chest wall	Costochondritis	Breast disease
	Chest wall trauma or inflammation	Rib fracture
	Herpes zoster (shingles)	Musculoskeletal injury/spasm

4.3.1. Evaluation of Acute Chest Pain With Suspected Gastrointestinal Syndromes

▶

1. In patients with recurrent acute chest pain without evidence of a cardiac or pulmonary cause, evaluation for gastrointestinal causes is reasonable. (*IIa, C-LD*)

4.3.2. Evaluation of Acute Chest Pain With Suspected Anxiety and Other Psychosomatic Considerations



1. For patients with recurrent, similar presentations for acute chest pain with no evidence of a physiological cause on prior diagnostic evaluation including a negative workup for myocardial ischemia, referral to a cognitive-behavioral therapist is reasonable. (*IIa, B-R*)

4.3.3. Evaluation of Acute Chest Pain in Patients With Sickle Cell Disease



1. In patients with sickle cell disease who report acute chest pain, emergency transfer by EMS to an acute care setting is recommended. (*I, B-NR*)



2. In patients with sickle cell disease who report acute chest pain, ACS should be excluded. (*I, C-LD*)

Figure 11. Pretest Probabilities of Obstructive CAD in Symptomatic Patients According to Age, Sex, and Symptoms

Pretest Probabilities of Obstructive CAD in Symptomatic Patients
 (A) according to age, sex, and symptoms;
 (B) according to age, sex, symptoms, and CAC

Age, y	Chest Pain		Dyspnea	
	Men	Women	Men	Women
30–39	≤4	≤5	0	3
40–49	≤22	≤10	12	3
50–59	≤32	≤13	20	9
60–69	≤44	≤16	27	14
70+	≤52	≤27	32	12

A Pretest probability based on age, sex, and symptoms

Low ≤15%	Intermediate–High >15%
-------------	---------------------------

B Pretest probability based on age,

sex, symptoms, and CAC score⁺

≤15%	>15%–50%	>50%
CAC 1–99	CAC ≥100–999	CAC ≥1,000

The Pretest Probability shown is for patients with anginal symptoms. Patients with lower risk symptoms would be expected to have lower PTP.

The darker green and orange shaded regions denote the groups in which non-invasive testing is most beneficial (pre-test probability >15%). The light green shaded regions denote the groups with pre-test probability of CAD ≤15% in which the testing for diagnosis may be considered based on clinical judgement.

If CAC available, can use to estimate pretest probability based on CAC Score.

Adapted and modified from Juarez-Orozco *ESC* 2019 Nov 1;20(11):1198-1207.

+ Winther, S. et al. *J Am Coll Cardiol.* 2020 Nov 24;76(21):2421-2432.

5.1.2. Low-Risk Patients With Stable Chest Pain and No Known CAD



1. For patients with stable chest pain and no known CAD presenting to the outpatient clinic, a model to estimate pretest probability of obstructive CAD is effective to identify patients at low risk for obstructive CAD and favorable prognosis in whom additional diagnostic testing can be deferred. (*I, B-NR*)



2. For patients with stable chest pain and no known CAD categorized as low risk, CAC testing is reasonable as a first-line test for excluding calcified plaque and identifying patients with a low likelihood of obstructive CAD. (*IIa, B-R*)



3. For patients with stable chest pain and no known CAD categorized as low risk, exercise testing without imaging is reasonable as a first-line test for excluding myocardial ischemia and determining functional capacity in patients with an interpretable ECG. (*IIa, B-NR*)

5.1.3. Intermediate-High Risk Patients With Stable Chest Pain and No Known CAD

Anatomic Testing



1. For intermediate-high risk patients with stable chest pain and no known CAD, CCTA is effective for diagnosis of CAD, for risk stratification, and for guiding treatment decisions. (*I, A*)

Stress Testing



2. For intermediate-high risk patients with stable chest pain and no known CAD, stress imaging (stress echocardiography, PET/SPECT MPI or CMR) is effective for diagnosis of myocardial ischemia and for estimating risk of MACE. (*I, B-R*)



3. For intermediate-high risk patients with stable chest pain and no known CAD for whom rest/stress nuclear MPI is selected, PET is reasonable in preference to SPECT, if available to improve diagnostic accuracy and decrease the rate of nondiagnostic test results. (*IIa, B-R*)



4. For intermediate-high risk patients with stable chest pain and no known CAD with an interpretable ECG and ability to achieve maximal levels of exercise (≥5 METs), exercise electrocardiography is reasonable. (*IIa, B-R*)

- ▶ 5. In intermediate-high risk patients with stable chest pain selected for stress MPI using SPECT, the use of attenuation correction or prone imaging may be reasonable to decrease the rate of false-positive findings. (*IIb, B-NR*)

Assessment of Left Ventricular Function


- ▶ 6. In intermediate-high risk patients with stable chest pain who have pathological Q waves, symptoms or signs suggestive of heart failure, complex ventricular arrhythmias, or a heart murmur with unclear diagnosis, use of TTE is effective for diagnosis of resting left ventricular systolic and diastolic ventricular function and detection of myocardial, valvular, and pericardial abnormalities. (*I, B-NR*)

Secondary Diagnostic Testing: What to Do If Index Test Results Are Positive or Inconclusive

SEQUENTIAL OR ADD-ON DIAGNOSTIC TESTING

- ▶ 7. For intermediate-high risk patients with stable chest pain and known coronary stenosis of 40% to 90% in a proximal or middle coronary segment on CCTA, FFR-CT can be useful for diagnosis of vessel-specific ischemia and to guide decision-making regarding the use of coronary revascularization. (*IIa, B-NR*)
- ▶ 8. For intermediate-high risk patients with stable chest pain after an inconclusive or abnormal exercise ECG or stress imaging study, CCTA is reasonable. (*IIa, B-NR*)
- ▶ 9. For intermediate-high risk patients with stable chest pain and no known CAD undergoing stress testing, the addition of CAC testing can be useful. (*IIa, B-NR*)
- ▶ 10. For intermediate-high risk patients with stable chest pain after inconclusive CCTA, stress imaging is reasonable. (*IIa, B-NR*)
- ▶ 11. For intermediate-high risk patients with stable chest pain after a negative stress test but with high clinical suspicion of CAD, CCTA or ICA may be reasonable. (*IIb, C-EO*)

Figure 12. Clinical Decision Pathway for Patients With Stable Chest Pain and No Known CAD



5.2. Patients With Known CAD Presenting With Stable Chest Pain COR LOE

- ▶ 1. For patients with obstructive CAD and stable chest pain, it is recommended to optimize GDMT. (*I, A*)
- ▶ 2. For patients with known nonobstructive CAD and stable chest pain, it is recommended to optimize preventive therapies. (*I, C-EO*)

5.2.1. Patients With Obstructive CAD Who Present With Stable Chest Pain


Index Diagnostic Testing

ANATOMIC TESTING

- ▶ 1. For patients with obstructive CAD who have stable chest pain despite GDMT and moderate-severe ischemia, ICA is recommended for guiding therapeutic decision-making. (*I, A*)
- ▶ 2. For patients with obstructive CAD who have stable chest pain despite optimal GDMT, those referred for ICA without prior stress testing benefit from FFR or instantaneous wave free ratio. (*I, A*)
- ▶ 3. For symptomatic patients with obstructive CAD who have stable chest pain with CCTA-defined ≥50% stenosis in the left main coronary artery, obstructive CAD with FFR with CT ≤0.80, or severe stenosis (≥70%) in all 3 main vessels, ICA is effective for guiding therapeutic decision-making. (*I, B-R*)
- ▶ 4. For patients who have stable chest pain with previous coronary revascularization, CCTA is reasonable to evaluate bypass graft or stent patency (for stents ≥3 mm). (*Ila, B-NR*)

- ▶ 5. For patients with obstructive CAD who have stable chest pain despite optimal GDMT, stress PET/SPECT MPI, CMR, or echocardiography is recommended for diagnosis of myocardial ischemia, estimating risk of MACE, and guiding therapeutic decision-making. **(I, B-NR)**
- ▶ 6. For patients with obstructive CAD who have stable chest pain despite optimal GDMT, when selected for rest/stress nuclear MPI, PET is reasonable in preference to SPECT, if available, to improve diagnostic accuracy and decrease the rate of nondiagnostic test results. **(IIa, B-R)**
- ▶ 7. For patients with obstructive CAD who have stable chest pain despite GDMT, exercise treadmill testing can be useful to determine if the symptoms are consistent with angina pectoris, assess the severity of symptoms, evaluate functional capacity and select management, including cardiac rehabilitation. **(IIa, B-R)**
- ▶ 8. For patients with obstructive CAD who have stable chest pain symptoms undergoing stress PET MPI or stress CMR, the addition of MBFR is useful to improve diagnosis accuracy and enhance risk stratification. **(IIa, B-NR)**

Figure 13. Clinical Decision Pathway for Patients With Stable Chest Pain (or Equivalent) Symptoms With Prior MI, Prior Revascularization, or Known CAD on Invasive Coronary Angiography or CCTA, Including Those With Nonobstructive CAD



5.2.1.1. Patients With Prior Coronary Artery Bypass Surgery With Stable Chest Pain

- ▶ 1. In patients who have had prior coronary artery bypass surgery presenting with stable chest pain whose noninvasive stress test results show moderate to severe ischemia, or in those suspected to have myocardial ischemia with indeterminate/nondiagnostic stress test, ICA is recommended for guiding therapeutic decision-making. **(I, C-LD)**
- ▶ 2. In patients who have had prior coronary artery bypass surgery presenting with stable chest pain who are suspected to have myocardial ischemia, it is reasonable to perform stress imaging or CCTA to evaluate for myocardial ischemia or graft stenosis or occlusion. **(IIa, C-LD)**

5.2.2. Patients With Known Nonobstructive CAD Presenting With Stable Chest Pain

Index Diagnostic Testing: Selecting the Appropriate Test

ANATOMIC TESTING

- ▶
 1. For symptomatic patients with known nonobstructive CAD who have stable chest pain, CCTA is reasonable for determining atherosclerotic plaque burden and progression to obstructive CAD, and guiding therapeutic decision-making. (*Ila, B-NR*)

- ▶
 2. For patients with known coronary stenosis from 40% to 90% on CCTA, FFR can be useful for diagnosis of vessel-specific ischemia and to guide decision-making regarding the use of ICA. (*Ila, B-NR*)

STRESS TESTING

- ▶
 3. For patients with known extensive nonobstructive CAD with stable chest pain symptoms, stress imaging (PET/SPECT, CMR, or echocardiography) is reasonable for the diagnosis of myocardial ischemia. (*Ila, C-LD*)

5.2.3. Patients With Ischemia and No Obstructive CAD (INOCA)


- ▶
 1. For patients with persistent stable chest pain and nonobstructive CAD and at least mild myocardial ischemia on imaging, it is reasonable to consider invasive coronary function testing to improve the diagnosis of coronary microvascular dysfunction and to enhance risk stratification. (*Ila, B-NR*)

- ▶
 2. For patients with persistent stable chest pain and nonobstructive CAD, stress PET MPI with MBFR is reasonable to diagnose microvascular dysfunction and enhance risk stratification. (*Ila, B-NR*)

- ▶
 3. For patients with persistent stable chest pain and nonobstructive CAD, stress CMR with the addition of MBFR measurement is reasonable to improve diagnosis of coronary myocardial dysfunction and for estimating risk of MACE. (*Ila, B-NR*)

- ▶
 4. For patients with persistent stable chest pain and nonobstructive CAD, stress echocardiography with the addition of coronary flow velocity reserve measurement may be reasonable to improve diagnosis of coronary myocardial dysfunction and for estimating risk of MACE. (*Ilb, C-EO*)

Figure 14. Clinical Decision Pathway for INOCA



Class of Recommendations and Level of Evidence

Class (Strength) of Recommendation

CLASS I (STRONG)

Benefit >>> Risk

Suggested phrases for writing recommendations:

- Is recommended
- Is indicated/useful/effective/beneficial
- Should be performed/administered/other
- Comparative-Effectiveness Phrases[†]:
 - Treatment/strategy A is recommended/indicated in preference to treatment B
 - Treatment A should be chosen over treatment B

CLASS IIa (MODERATE)

Benefit >> Risk

Suggested phrases for writing recommendations:

- Is reasonable
- Can be useful/effective/beneficial
- Comparative-Effectiveness Phrases[†]:
 - Treatment/strategy A is probably recommended/indicated in preference to treatment B
 - It is reasonable to choose treatment A over treatment B

CLASS IIb (WEAK)

Benefit \geq Risk

Suggested phrases for writing recommendations:

- May/might be reasonable
- May/might be considered
- Usefulness/effectiveness is unknown/unclear/uncertain or not well established

CLASS III: No Benefit (MODERATE)
(Generally, LOE A or B use only)

Benefit = Risk

Suggested phrases for writing recommendations:

- Is not recommended
- Is not indicated/useful/effective/beneficial
- Should not be performed/administered/other

CLASS III: Harm (STRONG)

Risk > Benefit

Suggested phrases for writing recommendations:

- Potentially harmful
- Causes harm
- Associated with excess morbidity/mortality
- Should not be performed/administered/other

Level (Quality) of Evidence[‡]

LEVEL A

- High-quality evidence[‡] from more than 1 RCT
- Meta-analyses of high-quality RCTs
- One or more RCTs corroborated by high-quality registry studies

LEVEL B-R

(Randomized)

- Moderate-quality evidence[‡] from 1 or more RCTs
- Meta-analyses of moderate-quality RCTs

LEVEL B-NR

(Nonrandomized)

- Moderate-quality evidence[‡] from 1 or more well-designed, well-executed nonrandomized studies, observational studies, or registry studies
- Meta-analyses of such studies

LEVEL C-LD

(Limited Data)

- Randomized or nonrandomized observational or registry studies with limitations of design or execution
- Meta-analyses of such studies
- Physiological or mechanistic studies in human subjects

LEVEL C-EO

(Expert Opinion)

- Consensus of expert opinion based on clinical experience

COR and LOE are determined independently (any COR may be paired with any LOE).

A recommendation with LOE C does not imply that the recommendation is weak. Many important clinical questions addressed in guidelines do not lend themselves to clinical trials. Although RCTs are unavailable, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

* The outcome or result of the intervention should be specified (an improved clinical outcome or increased diagnostic accuracy or incremental prognostic information).

† For comparative-effectiveness recommendations (COR I and IIa; LOE A and B only), studies that support the use of comparator verbs should involve direct comparisons of the treatments or strategies being evaluated.

‡ The method of assessing quality is evolving, including the application of standardized, widely used, and preferably validated evidence grading tools; and for systematic reviews, the incorporation of an Evidence Review Committee.

COR indicates Class of Recommendation; EO, expert opinion; LD, limited data; LOE, Level of Evidence; NR, nonrandomized; R, randomized; RCT, randomized controlled trial.

Source Citation

Disclaimer

Find...



TOOLS

[Log In to use Notes](#)

A+ Increase Font Size

A- Decrease Font Size



FILTERS

[Clear All](#)

[Level](#)

[Class](#)

[Inclusion Criteria](#)

[Recommendation Scope](#)

[SHOP](#)

[All Products](#)

[GUIDELINES LIBRARY](#)

[Pocket Guides](#)

[ABOUT](#)

[About Guideline Central](#)

[Careers](#)

[Guideline Central Blog](#)

[Frequently Asked Questions](#)

[CONTACT US](#)

[Contact Form](#)

[CART](#)

[FEEDBACK](#)

